

Research article

Indoor-outdoor concentrations of fine particulate matter in school building microenvironments near a mine tailing deposit

Leonardo Martínez^{1,2}, Stephanie Mesías Monsalve³, Karla Yohannessen Vásquez³, Sergio Alvarado Orellana^{3,5,6}, José Klarián Vergara⁷, Miguel Martín Mateo^{5,8}, Rogelio Costilla Salazar⁹, Mauricio Fuentes Alburquerque⁴, Ana Maldonado Alcaíno³, Rodrigo Torres^{3,4}, and Dante D. Cáceres Lillo^{3,5,6,*}

¹ University of Georgia, School of Public Health, Department of Epidemiology and Biostatistics, Athens, Georgia, USA

² Department of Preventive Medicine, Icahn School of Medicine, Mount Sinai, New York, New York, USA

³ Programa de Salud Ambiental, Escuela de Salud Pública, Facultad de Medicina, Universidad de Chile, Santiago, Chile

⁴ Programa de Bioestadística, Escuela de Salud Pública, Facultad de Medicina, Universidad de Chile, Santiago, Chile

⁵ Grups de Recerca d'Amèrica i Àfrica Llatines GRAAL, Unitat de Bioestadística, Facultat de Medicina, Universitat Autònoma de Barcelona, Barcelona, Spain

⁶ Facultad de Ciencias de la Salud, Universidad de Tarapacá, Arica, Chile

⁷ Departamento de Prevención de Riesgos y Medioambiente, Universidad Tecnológica Metropolitana, Santiago, Chile

⁸ Facultad de Ciencias Médicas, de la Salud y la Vida, Universidad Internacional del Ecuador, Ecuador

⁹ División de Ciencias de la Vida, Campus Irapuato-Salamanca, Universidad de Guanajuato, Guanajuato, México

*** Correspondence:** Email: dcaceres@med.uchile.cl; Tel: +56 2 9786539.

Abstract: Indoor air quality in school classrooms is a major pediatric health concern because children are highly susceptible to adverse effects from xenobiotic exposure. Fine particulate matter (PM_{2.5}) emitted from mining waste deposits within and near cities in northern Chile is a serious environmental problem. We measured PM_{2.5} in school microenvironments in urban areas of Chañaral, a coastal community whose bay is contaminated with mine tailings. PM_{2.5} levels were measured in

six indoor and outdoor school environments during the summer and winter of 2012 and 2013. Measurements were taken during school hours on two consecutive days. Indoor PM_{2.5} concentrations were 12.53–72.38 $\mu\text{g}/\text{m}^3$ in the summer and 21.85–100.53 $\mu\text{g}/\text{m}^3$ in winter, while outdoor concentrations were 11.86–181.73 $\mu\text{g}/\text{m}^3$ in the summer and 21.50–93.07 $\mu\text{g}/\text{m}^3$ in winter. Indoor/outdoor ratios were 0.17–2.76 in the summer and 0.64–4.49 in winter. PM_{2.5} levels were higher in indoor microenvironments during the winter, at times exceeding national and international recommendations. Our results demonstrate that indoor air quality Chañaral school microenvironments is closely associated with outdoor air pollution attributable to the nearby mine tailings. Policymakers should enact environmental management strategies to minimize further environmental damage and mitigate the risks that this pollution poses for pediatric health.

Keywords: particulate matter; schools; mine tailings

1. Introduction

Air quality in the school environment that has taken on increasing public health importance in recent years. Several studies have reported that school indoor microenvironments are among the most highly contaminated sites, finding high levels of particles, gases, and microorganisms associated with acute and chronic health problems [1-6]. A correlation exists between the concentration of pollutants and health problems in school children. These observed health effects are predominantly respiratory, including asthma, allergies, respiratory infections, chronic obstructive pulmonary disease, and the risk for developing other pulmonary obstructive diseases in adulthood [5,7,8]. These findings are concerning, given that children typically spend 6–10 hours per day in elementary schools, preschools, or nurseries [5,9]. Children are especially susceptible to the adverse effects of air pollution, due to their immature physiological and metabolic systems, their immunological defenses are not fully developed [9-11]. Additionally, children's normal activity patterns tend to stir up dust, increasing exposure to pollutants [12,13]. On the other hand, because of their smaller stature, children are exposed to a higher concentration of inhaled aerosol particles. Because their airways are smaller, children inhale a higher level of air per unit body weight compared to adults. Smaller air particles, such as PM_{2.5} and ultrafine particles, are considered more harmful than larger particles [14-16]. In addition to size, the chemical composition of PM can determine its dangerousness while components such as transition metals (iron, copper, nickel, and chromium) are known to produce severe toxicity on exposed tissues [17-19]. Reactive oxygen species generated during metabolism of air contaminants are responsible for inflammatory processes that occur in response to air pollutant exposure [20].

Mining has been Chile's main economic activity for decades, operating primarily from central and northern regions of the country [21]. Mining provides economic benefits but also creates environmental damage, due to residues left after mineral processing. One such environmental liability is mine tailings of ore processing, containing high levels of toxic metals [22-24].

Chañaral is a coastal city in northern Chile whose main commercial activity is mining and fishing. During the last century, it had a great mining boom which resulted in large beneficial to the country. However, this activity generated a lot of waste that ended up in the Chañaral's bay, embanking and polluting the beach. This massive pollution of the coastline has affected the marine

flora and fauna [25-27]. This phenomenon of coastal mining waste pollution has been described in different places of the world [28-31].

In 2011, environmental Chilean authorities identified at least 603 mine tailing deposits, many of which are adjacent to highly-populated areas [32]. The bay of Chañaral in northern Chile is one such area where strong prevailing winds carry dust from the beach towards urban areas during spring and summer [33,34]. The aim of this study is to determine fine particulate matter concentrations in school microenvironments settings and to measure the relationship between indoor and outdoor levels.

2. Materials and Methods

2.1. Study design and schools sampling

We performed a cross-sectional study of a sampling frame of 10 schools in Chañaral, from which we randomly selected six schools distributed throughout the city (Figure 1). Sets of measurements were performed during the summer (December 2012) and repeated during the winter (July 2013). The microenvironments studied were classrooms, offices, and playgrounds on school property. Administrative offices and classrooms were classified as indoor environments and playgrounds as outdoor environments. The characteristics of the sampling areas in preschools (PS) and elementary schools (ES) are described in Table 1.



Figure 1. Map of Chañaral, Chile and locations of the participating schools. PS, preschools; ES, Elementary schools.

Table 1. Characteristics of preschool and elementary school sampling areas, Atacama Region, Chile.

ID	Room ventilation	Office ventilation	Indoor floor type	Outdoor floor type	Location(s)	Comments
ES1	Small iron windows to outside closed throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Concrete/Ground	Hilltop, near railway line	Elementary school located near train line, farther from beach than most of the other schools. Daily cleanup after school hours
ES2	Small aluminum windows to outside open throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Concrete	On main two-way street	School located farther from beach than most of the other schools. Daily cleanup after school hours
ES3	Aluminum windows to outside closed throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Concrete/Ground	Near Main Street	Playground remodeling construction activities occurred throughout the process. Daily cleanup after school hours
PS1	Wood windows to outside closed throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Ground	On Main Street and adjacent to beach	School located near the beach, adjacent to the municipal greenhouse. Daily cleanup after school hours
PS2	Small aluminum windows to outside open throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Concrete	On Main Street and adjacent to beach	School located near the beach, with playground adjacent to a bus station. Daily cleanup after school hours
PS3	Small aluminum windows to outside open throughout the process	One aluminum window to exterior, closed throughout the process	Vinyl	Concrete/Ground	On main two-way street	School located near the beach, in an obviously impacted area, and adjacent to a local mini-zoo. Daily cleanup after school hours

PS: Preschool; ES: Elementary school

2.2. Study area

Chañaral is a northern coastal city with a warm desert climate, abundant clouds, and scarce rain. Chañaral has a narrow thermal amplitude with low contrast, both across seasons and between night and day, and prevailing west-east winds [35]. The ocean currents stir up the tailing mud when the coast is soaked by high tides [26,34].

2.3. *PM_{2.5} indoor vs outdoor gravimetric sampling and meteorological parameters*

PM_{2.5} mass concentration was measured by gravimetric analysis in Chester LabNet Laboratory, (www.chesterlab.net), a National Environmental Laboratory Accreditation Conference (NELAC) accredited laboratory (Tigard, OR, USA). Personal environmental monitoring system (PEMS)[®] equipment model 761-203A, SKC, Eighty Four, PA) containing 37 mm pre-weighed Teflon filters was used to collect PM_{2.5}, using a 44XR Universal Sample Pump (SKC Inc, Eight Four, Pennsylvania, USA). A flow of 4 L/min was set, and variation was monitored daily by means of a soap bubble electronic SKC ULTRAFLO[®] calibrator. PM_{2.5} was measured in indoor and outdoor microenvironments simultaneously. All measurements were performed during school hours on two consecutive days, for eight hours per day, resulting in a total measurement time of 16 hours for each school per season. In indoor environments, particle collectors were placed at a height of one meter and a distance of 50 cm from the wall, in a corner opposite the main windows. In outdoor areas, collectors were placed on the playground, also at a height of one meter. A total of 36 measurements were performed at each site.

Meteorological station equipment with adequate coverage of the target area was used to record wind velocity, temperature, relative humidity, solar radiation, barometric pressure, and precipitation. Meteorological parameters were measured according to established guidelines [36].

2.4. *Data analysis*

Descriptive and exploratory analyses were performed on the data collected. To compare PM_{2.5} mass concentration between stations and microenvironments two-sample Wilcoxon rank-sum test was used. We also analyzed indoor-outdoor ratios (I/O), with the assumption that I/O values higher than unity (1) would suggest another source of emissions into the studied areas and/or outside infiltration [37,38]. To evaluate indoor and outdoor relationships among offices, classrooms and playground during the sampling period Spearman's correlation coefficient was used, respectively. Statistical analysis was performed using STATA 11.1 software.

2.5. *Ethical considerations*

The study was conducted as part of FONIS project SA11|2224 and was approved by the ethical committee of the University of Chile School of Medicine [39].

3. Results

3.1. *PM_{2.5} levels and indoor-outdoor relationship in schools' microenvironments*

Figure 2 shows PM_{2.5} mass concentrations by season and microenvironment. During summer, PM_{2.5} mass concentrations were 12.53–72.38 $\mu\text{g}/\text{m}^3$ within indoor microenvironments and 11.86–181.73 $\mu\text{g}/\text{m}^3$ in outdoor microenvironments. In winter, PM_{2.5} concentrations were 21.5–93.1 and 21.9–100.5 $\mu\text{g}/\text{m}^3$ in outdoor and indoor settings, respectively. No significant differences for indoor and outdoor PM_{2.5} median concentrations for summer ($p = 0.6199$) and winter ($p = 0.2026$) were observed, respectively. In summer, the highest indoor value was found in ES2 (office) and the highest outdoor values in PS1 and PS2 playgrounds, respectively. The highest values in winter were

found in PS1 (all microenvironments), and PS2, ES3 for classroom, respectively (see supplementary data).




Figure 2. Summer and winter PM_{2.5} concentration and indoor/outdoor ratio in school microenvironments in Chañaral, Chile. PS, preschools; ES, Elementary schools.

When comparing the median concentrations of PM_{2.5} between summer and winter season, not statistically significant differences for offices ($p = 0.3367$) and playgrounds ($p = 0.4233$) were observed, respectively. However, the median concentration of PM_{2.5} in classrooms in winter period was significantly higher compared with summer ($p = 0.0104$).

Table 2 shows indoor vs outdoor PM_{2.5} relationship by season and microenvironment. Summer I/O ratios for offices and classrooms versus playgrounds varied from 0.17 to 2.76 and 0.12 to 2.23, respectively. The relationship between PM_{2.5} levels for offices and classrooms for ES1, PS1, PS2 schools compared with playground was less than unity (1), indicating that during the period sampled, the outdoor levels were higher. In some cases that outdoor concentrations were nearly three to eight times higher than indoor concentrations. This can be explained by any of the following reasons; in ES1 during sampling period, there were works of removal of construction debris outside school, this generated locally resuspension of dust that may have affected the measurement. For PS1 and PS2, it can be explained by its proximity to the area of the deposit of tailings on the beach. In addition, another factor was that the floor of these schools was not completely paved, at least 50% were soil and sand. To ES2 and PS3 I/O relationship was greater than unity (1), indicating higher levels of PM_{2.5} in the microenvironments of offices and classrooms compared with playground. In this preschool, the playground was paved, but there was unpaved area near to the classrooms gate, where it would be entering the dust, which is carried on their clothes and shoes for children and teachers during the school day.

During winter, I/O ratios were 0.17–2.76 for offices vs. playgrounds and 0.12–2.23 for classrooms vs. playgrounds. To classroom vs playground the I/O was greater than 1.0 except in PS3, indicating a higher concentration of PM_{2.5} in the classroom compared with playground. This concentration was up to 4.49 times the outdoor as shown in ES3 (Table 2). This school, the classroom and offices are adjacent to an unpaved area, which could explain these high levels of PM_{2.5}. On the other hand, the relationship between offices and playgrounds was around unity (1.0).

Correlation analysis during the sampling period showed no clear relationship between outdoor-indoor PM_{2.5} levels for playground versus classrooms ($r = 0.23$; $p = 0.4705$). Instead, the

correlation of PM_{2.5} levels in playground versus offices was moderate, near of statistical significance ($r = 0.56$; $p = 0.0548$).

Table 2. Indoor/outdoor ratio to PM2.5 concentration in school microenvironments. Chañaral, Chile.

School	Summer		Winter	
	Indoor/Outdoor		Indoor/Outdoor	
	Office/Play	Classroom/Play	Office/Play	Classroom/Play
ES1	0.71	0.72	0.82	1.14
ES2	2.76	1.05	1.44	1.21
ES3	1.38	1.04	1.69	4.49
PS1	0.17	0.12	1.00	1.08
PS2	0.38	0.16	0.86	2.24
PS3	1.27	2.23	0.64	0.97
Median	1.04	0.78	0.93	1.67

ES: Elementary Schools; PS: Preschools; Play: Playground.

3.2. Meteorological parameters

Figure 3 demonstrates the direction of the local wind patterns in Chañaral as well as the wind velocity. During 42–56% of the period studied, the prevailing wind direction was West-Northwest (November–March, which are the spring and summer months in South America). During fall and winter (April–June), the wind direction was west, northeast, and east 27, 15 and 20% of the time, respectively (Figure 3a). Wind velocity was more variable during the spring and summer months (December–February) as compared to the fall and winter months (Figure 3b). During this period, 75% of measurements exceeded 4 m/s, and the strongest winds generally occurred between 10:00 and 18:00 hours (see supplementary data).

4. Discussion

This study aimed to determine the concentrations of fine particulate matter (PM_{2.5}) at school microenvironments in two different seasons in the coastal city of Chañaral in northern Chile, which bay is heavily polluted with ore waste of copper mines.

The ratio of indoor versus outdoor concentration levels varied by season and were generally high during the winter season. During the summer, there were not significant differences in the median concentrations in indoor compared to outdoor microenvironments. However, concentrations of the two schools closest to the coastline (both preschool level elementary schools) had high concentrations of PM_{2.5}. We also did not observe a clear difference in indoor microenvironments. In winter, the median PM_{2.5} concentrations were similar in offices and patios, however, levels in classrooms were higher than in playgrounds. The very high levels seen in outdoor environments during the summer may be associated with atmospheric conditions prevalent during the days we monitored the schools. This weather phenomenon may have influenced the concentrations of particulate matter and dust in suspension in the study area. Some authors have reported that




Figure 3. Wind characteristics, November 2013 to May 2013, Chañaral, Chile.

particulate matter concentration is modulated by wind speed, observing lower concentrations for wind speeds under 2 m/s and higher concentrations for stronger winds [40]. Mining waste is amongst the largest generators of dust that pollute the atmosphere. Pollution in coastal areas includes a complex mixture of tailings containing metal, sand, and the effect of sea water, generating particulate matter carried by the wind from surrounding urban areas and communities [24,26,41-44].

This finding may explain the higher particulate matter levels observed in summer, when the wind speed was significantly higher [45,46]. Other authors have reported a similar relationship between wind speed and concentration, albeit with slightly different values, finding, for example, that PM concentrations were higher when wind speeds were above 4 m/s [46-48]. During this study, the 75th percentile of wind velocity measurements was about 4 m/s, with higher speeds in summer vs. winter (Figure 3b). During the study period, no rainfall was recorded. The average relative humidity

for the period was 70% (range 34.3–89.9). Barometric pressure and solar radiation were practically constant over the period. Neary and Garcia-Chevesich reported that during spring and summer, local prevailing winds blow towards the coast in this region (west to east), carrying dust from the bay and moving particles towards the urban area of Chañaral [33].

High indoor particulate matter levels in preschools may be attributable to the schools' proximity to the bay polluted with mine tailing deposits, as well as the activity of children in classrooms, which can resuspend particulate matter. During the summer, the preschools closest to the beach kept their windows closed during school hours to prevent dust from entering school offices and classes. This method was relied upon particularly in schools lacking formal ventilation. In the winter, the windows also remained closed most of the time, in this case to keep the classrooms warm, especially in the classrooms where the children spent the most time. The high indoor particular matter levels observed in our study could be explained by infiltration from the outdoor environment and the movement of children during the school day. As it is seen in summer time distribution of PM_{2.5} concentration levels, increase, which would be associated with periods of recess and lunchtime. During these periods of time, children are moving, causing particle resuspension. In preschools (PS), a greater variation in levels of PM compared to basic schools (winter and summer), which could be related to the proximity to the bay were observed. Moreover, the elementary school (ES2) is at higher altitudes and farther from the area contaminated beach.

Seasonal differences were not as apparent in the offices as in the classrooms, suggesting that the variations in classroom levels were attributable to children's activity during school hours. The administrative offices where PM_{2.5} samplers were installed, there were no specific sources of emission of particles, such as photocopying or other equipment. The offices were occupied by school principals and teachers.

Various authors have noted that indoor particulate matter levels are closely related to the presence and activity of students, as well as to levels in outdoor environments [23,49-51]. Several international studies have found results similar to those reported here. Studies performed in England and the Netherlands reported indoor PM concentrations of 19.0–30.0 and 7.7–52.8 $\mu\text{g}/\text{m}^3$, respectively [52,53]. A study by Diapouli et al., which measured PM_{2.5} levels during winter in seven schools, reported indoor concentrations of 22.1–40.3 $\mu\text{g}/\text{m}^3$ and outdoor concentrations of 38.6–99.0 $\mu\text{g}/\text{m}^3$, similar to the values observed in Chañaral (52.7 and 30.9 $\mu\text{g}/\text{m}^3$, respectively) [54]. However, another comparable study by Madureira et al., measuring PM_{2.5} concentration at 11 schools in Porto, Portugal, reported concentrations of 95 $\mu\text{g}/\text{m}^3$ indoors and 115 $\mu\text{g}/\text{m}^3$ outdoors; these values were significantly higher than those measured in Chañaral [55]. A study of six elementary schools in Iran (2013) reported average indoor levels of 29.1–69.1 $\mu\text{g}/\text{m}^3$ during the school day, similar to our results [56]. It is important to note that in the studies described above, the main source of particulate matter was vehicular traffic, a factor that is likely marginal in Chañaral.

Nkosi et al. (2015) conducted a study in 11 South Africa communities, reported an association between proximity to deposits of mine waste and effects on the respiratory health of children and adolescents in 22 primary and secondary schools. The authors concluded the damaging effect of the mine on respiratory health and the proximity of the community to the mine dumps [57]. Studies on the composition of particulate matter in industrial areas of the Atacama region have shown to have a high content of heavy metals in PM₁₀ and PM_{2.5} [58-61], which may indicate the potential health risks of the exposed population, especially children, the elderly, and those with concurrent illness.

5. Conclusion

Our results revealed high levels of fine particulate matter during the summer and winter within Chañaral indoor school settings. The highest concentrations were in classrooms, particularly within preschools closest to the mining tailings. Indoor fine particulate matter levels exceeded Chilean and international recommendations, especially during winter months. Outdoor fine particulate matter concentrations also generally exceeded Chilean and international norms during both seasons studied. Children at these schools are exposed to poor air quality, whose main source is from particulate matter from a reservoir of mine tailings present at the Chañaral bay, putting children and other residents at high risk for many health problems. This is the first study of its kind in Chile, whose objective was to measure the concentration of particulate matter in school microenvironments in a polluted area with mine tailing. The composition of metals in particulate matter is urgently needed to clarify the risk of potential adverse effects on the health of the population.

Acknowledgement

We would like to thank the school administrators who allowed us to collect data at their facilities, the Department of Education and Department of the Environment, the Municipality of Chañaral neighborhood council, the field staff, the Fondo Nacional de Investigación y Desarrollo en Salud (FONIS N° SA11 / 2244), and the University of Chile School of Medicine, Public Health Department.

Conflict of interest

The authors have no conflicts of interest to declare.

Reference

1. Kim JL, Elfman L, Mi Y, et al. (2007) Indoor molds, bacteria, microbial volatile organic compounds and plasticizers in schools--associations with asthma and respiratory symptoms in pupils. *Indoor Air* 17: 153-163.
2. Hespel D, Qian J, Nazaroff WW, et al. (2012) Human occupancy as a source of indoor airborne bacteria. *PLoS One* 7: e34867.
3. Oeder S, Dietrich S, Weichenmeier I, et al. (2012) Toxicity and elemental composition of particulate matter from outdoor and indoor air of elementary schools in Munich, Germany. *Indoor Air* 22: 148-158.
4. Oeder S, Jorres RA, Weichenmeier I, et al. (2012) Airborne indoor particles from schools are more toxic than outdoor particles. *Am J Respir Cell Mol Biol* 47: 575-582.
5. Cartieaux E, Rzepka MA, Cuny D (2011) Indoor air quality in schools. *Arch Pediatr* 18: 789-796.
6. Abramson SL, Turner-Henson A, Anderson L, et al. (2006) Allergens in school settings: results of environmental assessments in 3 city school systems. *J Sch Health* 76: 246-249.
7. Rivas E, Barrios C S, Dorner P A, et al. (2008) Association between indoor contamination and respiratory diseases in children living in Temuco and Padre Las Casas, Chile. *Rev Méd Chile* 767-774.

8. Flores C, Solis M, Fortt A, et al. (2010) Sintomatología respiratoria y enfermedad pulmonar obstructiva crónica y su asociación a contaminación intradomiciliaria en el Área Metropolitana de Santiago: Estudio Platino. *Rev Chil Enferm Respir* 26: 72-80.
9. Zhang Q, Zhu Y (2012) Characterizing ultrafine particles and other air pollutants at five schools in South Texas. *Indoor Air* 22: 33-42.
10. Gilliland F, McConnell R, Peters J, et al. (1999) A theoretical basis for investigating ambient air pollution and children's respiratory health. *Environ Health Perspect* 107: 403-407.
11. Dockery D, Skerrett P, Walters D, et al. (2005.) Development of lung function. *Effects of air pollution on children's health and development : A review of the evidence*. Bonn, World Health Organization Special Programme on Health and Environment . European Centre for Environment and Health. 108-133.
12. Gavidia T, Brune MN, McCarty KM, et al. (2011) Children's environmental health--from knowledge to action. *Lancet* 377: 1134-1136.
13. Gavidia TG, Pronczuk de Garbino J, Sly PD (2009) Children's environmental health: an under-recognised area in paediatric health care. *BMC Pediatr* 9: 10.
14. Diociaiuti M, Balduzzi M, De Berardis B, et al. (2001) The two PM(2.5) (fine) and PM(2.5-10) (coarse) fractions: evidence of different biological activity. *Environ Res* 86: 254-262.
15. Li N, Hao M, Phalen RF, et al. (2003) Particulate air pollutants and asthma. A paradigm for the role of oxidative stress in PM-induced adverse health effects. *Clin Immunol* 109: 250-265.
16. Schwartz J, LM. N (2000) Fine particles are more strongly associated than coarse particles with acute respiratory health effects in school children. *Epidemiol* 11: 6-10.
17. Aust AE, Ball JC, Hu AA, et al. (2002) Particle characteristics responsible for effects on human lung epithelial cells. *Res Rep Health Eff Inst* 110: 1-65.
18. Gavett SH, Haykal-Coates N, Copeland LB, et al. (2003) Metal composition of ambient PM2.5 influences severity of allergic airways disease in mice. *Environ Health Perspect* 111: 1471-1477.
19. Okeson CD, Riley MR, Fernandez A, et al. (2003) Impact of the composition of combustion generated fine particles on epithelial cell toxicity: influences of metals on metabolism. *Chemosphere* 51: 1121-1128.
20. Peters J (2004) Epidemiologic investigation to identify chronic effects of ambient air pollutants in southern California. California Air Resources Board and the California No. 94-331 No. 94-331.
21. Méndez LM (1996) Historiografía minera de Chile (1870-1996). *Ensayo Bibliográfico. Dimensión Historica de Chile*. 67-89.
22. Csavina J, Field J, Taylor MP, et al. (2012) A review on the importance of metals and metalloids in atmospheric dust and aerosol from mining operations. *Sci Total Environ* 433: 58-73.
23. Jung MC (2008) Contamination by Cd, Cu, Pb, and Zn in mine wastes from abandoned metal mines classified as mineralization types in Korea. *Environ Geochem Health* 30: 205-217.
24. Meza-Figueroa D, Maier RM, de la OVM, et al. (2009) The impact of unconfined mine tailings in residential areas from a mining town in a semi-arid environment: Nacozari, Sonora, Mexico. *Chemosphere* 77: 140-147.
25. Lagos G, Velasco P (1999) Environmental Policies and Practices in Chilean Mining. In: Centre IDR, editor. *Mining and the environments. Cases studies from the Americas*. Canada: National Library of Canada.
26. Dold B (2006) Element Flows Associated with Marine Shore Mine Tailings Deposits. *Environ Sci Technol* 40: 752-758.

27. Lee MR, Correa JA, Castilla JC (2001) An assessment of the potential use of the nematode to copepod ratio in the monitoring of metals pollution. The Chanaral case. *Mar Pollut Bull* 42: 696-701.

28. Koski RA (2012) Metal Dispersion Resulting from Mining Activities in Coastal Environments: A Pathways Approach. *Oceanography* 25: 170-183.

29. Li M, Qi J, Zhang H, et al. (2011) Concentration and size distribution of bioaerosols in an outdoor environment in the Qingdao coastal region. *Sci Total Environ* 409: 3812-3819.

30. Martinez-Sanchez MJ, Navarro MC, Perez-Sirvent C, et al. (2008) Assessment of the mobility of metals in a mining-impacted coastal area (Spain, Western Mediterranean). *J Geochem Explor* 96: 171-182.

31. Violintzis C, Arditoglou A, Voutsas D (2009) Elemental composition of suspended particulate matter and sediments in the coastal environment of Thermaikos Bay, Greece: delineating the impact of inland waters and wastewaters. *J Hazard Mater* 166: 1250-1260.

32. SERNAGEOMIN (2015) Catastro Nacional de Depósitos de Relave. Depósitos Activos y No activos 2015 Servicio Nacional de Geología y Minería.

33. Neary D, Garcia-Chevesich P (2008) Hydrology and erosion impacts of mining derived coastal sand dunes, Chanaral Bay, Chile. *Hydrol Water Resour Arizona Southwest* 38: 47-52.

34. Vergara A (2011) Cuando el río suena, piedras trae: Relaves de cobre en la bahía de Chañaral, 1938-1990. *Cuadernos de historia Departamento de Ciencias Históricas, Universidad de Chile*: 135-151.

35. Juliá C, Montecinos S, Maldonado A (2008) Características Climáticas de la Región de Atacama. en Libro Rojo de la Flora Nativa y de los Sitios Prioritarios para su Conservación: Región de Atacama In: F.A. Squeo GAJRG, eds. , editor: Ediciones Universidad de la Serena, La Serena, Chile.

36. EPA (2006) Quality Assurance Handbook for Air Pollution Measurement Systems Volume IV: Meteorological Measurements. In: Program AAQM, editor. U.S. Environmental Protection Agency Office of Air Quality Planning and Standards Air Quality Assessment Division RTP, NC 27711.

37. Lim JM, Jeong JH, Lee JH, et al. (2011) The analysis of PM2.5 and associated elements and their indoor/outdoor pollution status in an urban area. *Indoor Air* 21: 145-155.

38. Massey D, Masih J, Kulshrestha A, et al. (2009) Indoor/outdoor relationship of fine particles less than 2.5 um(PM2.5) in residential homes locations in central Indian region. *Built Environ* 44: 2037-2045.

39. Yohannessen K, Alvarado S, Mesías S, et al. (2015) Exposure to fine particles by mine tailing and lung function effects in a panel of schoolchildren, Chañaral, Chile. *J Environ Prot* 6: 118-128.

40. Tian G, Fan SB, Huang YH, et al. (2008) Relationship between wind velocity and PM10 concentration & emission flux of fugitive dust source. *Huan Jing Ke Xue* 29: 2983-2986.

41. Moreno ME, Acosta-Saavedra LC, Meza-Figueroa D, et al. (2010) Biomonitoring of metal in children living in a mine tailings zone in Southern Mexico: A pilot study. *Int J Hyg Environ Health* 213: 252-258.

42. Hu H, Shine J, Wright RO (2007) The challenge posed to children's health by mixtures of toxic waste: the Tar Creek superfund site as a case-study. *Pediatr Clin North Am* 54: 155-175.

43. Ojelede M, Annegarn H, Kneen M (2012) Evaluation of aeolian emissions from gold mine tailings on the Witwatersrand. *Aeolian Res* 3: 477-486.

44. Bea SA, Ayora C, Carrera J, et al. (2010) Geochemical and environmental controls on the genesis of soluble efflorescent salts in coastal mine tailings deposits: a discussion based on reactive transport modeling. *J Contam Hydrol* 111: 65-82.

45. Stovern M, Betterton EA, Saez AE, et al. (2014) Modeling the emission, transport and deposition of contaminated dust from a mine tailing site. *Rev Environ Health* 29: 91-94.

46. Stovern M, Felix O, Csavina J, et al. (2014) Simulation of windblown dust transport from a mine tailings impoundment using a computational fluid dynamics model. *Aeolian Res* 14: 75-83.

47. Harrison RM, Yin J, Mark D, et al. (2001) Studies of coarse particle (2.5–10 μm) component in UK urban atmospheres. *Atmos Environ* 35: 3667-3679.

48. Alan J, Harrison RM, Baker J (2010) The wind speed dependence of the concentrations of airborne particulate matter and NOx. *Atmos Environ* 44: 1682-1690.

49. Lee S, Guo H, Li V, et al. (2002) inter-comparasion of air pollutant concentrations in different indoor environment in Hong Kong. *Atmos Environ* 36: 1929-1940.

50. John K, Karna S, Crist K, et al. (2007) Analysis of trace elements and ions in ambient fine particulate matter at three elementary schools in Ohio. *J Air Waste Manag Assoc* 57: 394-406.

51. Annesi-Maesano I, Moreau D, Caillaud D, et al. (2007) Residential proximity fine particles related to allergic sensitisation and asthma in primary school children. *Respir Med* 101: 1721-1729.

52. Janssen NAH, van Vliet PHN, Aarts F, et al. (2001) Assessment of exposure to traffic related air pollution of children attending schools near motorways. *Atmos Environ* 35: 3875-3884.

53. Wheeler AJ, Williams I, Beaumont RA, et al. (2000) Characterisation of Particulate Matter Sampled During a Study of Children's Personal Exposure to Airborne Particulate Matter in a UK Urban Environment. *Environ Monit Assess* 65: 69-77.

54. Diapouli E, Chaloulakou A, Mihalopoulos N, et al. (2008) Indoor and outdoor PM mass and number concentrations at schools in the Athens area. *Environ Monit Assess* 136: 13-20.

55. Madureira J, Paciencia I, Fernandes Ede O (2012) Levels and indoor-outdoor relationships of size-specific particulate matter in naturally ventilated Portuguese schools. *J Toxicol Environ Health A* 75: 1423-1436.

56. Mohammadyan M, Shabankhani B (2013) Indoor PM1, PM2.5, PM10 and outdoor PM2.5 concentrations in primary schools in Sari, Iran. *Arh Hig Rada Toksikol* 64: 371-377.

57. Nkosi V, Wichmann J, Voyi K (2015) Mine dumps, wheeze, asthma, and rhinoconjunctivitis among adolescents in South Africa: any association? *Int J Environ Health Res* 25: 583-600.

58. Jorquera H, Barraza F (2013) Source apportionment of PM10 and PM2.5 in a desert region in northern Chile. *Sci Total Environ* 444: 327-335.

59. Castilla JC (1983) Environmental impacts in sandy beaches of copper mine tailing at Chañaral, Chile. *Mar Pollut Bullet* 14: 159-464.

60. Parra S, Bravo MA, Quiroz W, et al. (2014) Distribution of trace elements in particle size fractions for contaminated soils by a copper smelting from different zones of the Puchuncavi Valley (Chile). *Chemosphere* 111: 513-521.

61. Jorquera H (2009) Source apportionment of PM10 and PM2.5 at Tocopilla, Chile (22 degrees 05' S, 70 degrees 12' W). *Environ Monit Assess* 153: 235-251.



AIMS Press

© 2016 Dante D. Cáceres Lillo et al., licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)