

AIMS Mathematics, 5(4): 4027–4044. DOI: 10.3934/math.2020259 Received: 17 October 2019 Accepted: 20 April 2020 Published: 27 April 2020

http://www.aimspress.com/journal/Math

# Research article

# New dark-bright soliton in the shallow water wave model

# Gulnur Yel<sup>1,\*</sup>, Haci Mehmet Baskonus<sup>2</sup> and Wei Gao<sup>2,3</sup>

- <sup>1</sup> Final International University, Kyrenia Mersin 10, Turkey
- <sup>2</sup> Harran University, Faculty of Education, Sanliurfa, Turkey
- <sup>3</sup> School of Information Science and Technology, Yunnan Normal University, Yunnan, China
- \* **Correspondence:** Email: gulnur.yel@final.edu.tr.

**Abstract:** In this paper, we employ the sine-Gordon expansion method to shallow water wave models which are Kadomtsev–Petviashvili–Benjamin–Bona–Mahony and the Benney–Luke equations. We construct many new complex combined dark-bright soliton, anti-kink soliton solutions for the governing models. The 2D, 3D and contour plots are given under the suitable coefficients. The obtained results show that the approach proposed for these completely integrable equations can be used effectively.

**Keywords:** the sine-Gordon expansion method; Kadomtsov–Petviashvili–Benjamin–Bona–Mahony equation; Benney–Luke equation; solitary wave solutions **Mathematics Subject Classification:** 35C08, 35Q51

# 1. Introduction

Solving completely integrable models is very important for observing and meaning dynamical behaviors of the physical systems. The shallow water wave model is one of the completely integrable models illustrating many physical problems. Euler is the first derived the water wave model in the 18th century. Using many models in applied mathematics and in physics such as Boussinesq equations, KdV equation, Davey–Stewarts on equation, Kadomtsev–Petviashvili (KP) equation, Benjamin–Bona–Mahony (BBM) equation are derived by Euler equations. The Benney–Luke equation is also water wave approximation to the Euler equation.

The KdV equation and the Benjamin–Bona–Mahony describe water waves that travel mainly in one direction. The Benney–Luke equation allows two-directional approximation to the Euler equations [1,2]. The KP equation is the 2D general form of KdV equation. It describes long

wavelength water waves in a weak dispersion medium. The Benjamin–Bona–Mahony (BBM) equation is also known regularized long-wave equation is an improvement form of KdV equation. It is derived to model long waves propagation in nonlinear dispersion. The KP equation and the BBM equations are used to symbolize for acoustic waves in fluids, long wavelength surface wave in liquids.

In this study we consider the combined the two equations which is KP-BBM equation given as [3];

$$\left(u_{t}+u_{x}-a\left(u^{2}\right)_{x}-bu_{xxt}\right)_{x}+ku_{yy}=0,$$
(1)

where a, b, k are constants and non-zero.

Many scientists have investigated the KP-BBM equation via various methods. In [4], the bifurcation of solitary waves of Eq. (1) is studied. The (3+1)-dimensional KP-BBM equation has been extracted to find analytical solutions by the auxiliary equation method [5]. Trigonometric and hyperbolic function solutions have been presented by new generalized (G'/G) expansion method [6]. The bell-shaped solitary wave and periodic wave solutions of KP-BBM equation have been obtained by bifurcation method and analyzed relation between of solutions under the different conditions [7]. The KP-BBM equation has been turned into Hirota bilinear form and obtained lump solutions by this way. The authors have investigated interactions lump-kink solutions and lump-soliton solutions [8].

In this work, we consider the Benney-Luke equation, the other model is derived from 3-dimensional Euler's equation.

The Benney-Luke equation is given by [9,10]

$$u_{tt} - u_{xx} + \alpha u_{xxxx} - \beta u_{xxtt} + u_t u_{xx} + 2u_x u_{xt} = 0,$$
(2)

where are arbitrary constants such that, is the Bond number capturing the effects of surface tension and gravity forces. Eq. (2) models an approximation of the full water wave equation and is formally suitable to describe the two way propagation of water waves in the presence of surface tension. Firstly, Eq. (2) is presented by Benney and Luke [11]. There are comprehensive studies such as numerical studies, stability analysis, traveling wave solutions, analytical solutions, Cauchy problem on this equation in the literature [1,9–15].

The shallow water wave equations used in geophysical fluid dynamics, oceanography, coastal engineering and atmospheric science. Especially, in the past fifty years, solitary waves and soliton solutions have been intensively studied in mathematical, physical and engineering fields. Some powerful methods are recommended to find solitary wave solutions such as Hirota's method [16], inverse scattering transform method [17], the solitary wave ansatz method [18,19], the tanh method [20], the extended auxiliary equation mapping, the extended direct algebraic mapping methods [21–33], Riemann–Hilbert method [34,35], the Lie symmetry method [36,37], and many others [38–43].

In this study, we focus on the sine-Gordon expansion method to construct to exact solutions of the KP-BBM equation and the Benney–Luke equation.

This paper rested as, the general properties of the mentioned method are given in Section 2. To obtain the soliton solutions the SGEM is applied to Eqs. (1) and (2), then 2D, 3D and contour simulations of the new solutions are plotted in Section 3. Finally, we give conclusion by submitted findings.

#### 2. The sine-Gordon expansion method

In this section, we describe SGEM. Firstly, we need to explain how we get two significant equations. We suppose that the sine-Gordon equation is given as following [44–47];

$$u_{xx} - u_{tt} = m^2 \sin(u), \tag{3}$$

Where u = u(x,t), *m* is a real constant. Applying the wave transform  $u = u(x,t) = U(\xi)$ ,  $\xi = x - ct$  to Eq. (3), we get the following nonlinear ordinary differential equation;

$$U'' = \frac{m^2}{\left(1 - c^2\right)} \sin\left(U\right),\tag{4}$$

Where  $U = U(\xi), \xi$  and *c* are the amplitude and velocity of the travelling waves, respectively. We integrate Eq. (4) and it can be written as follows;

$$\left[\left(\frac{U}{2}\right)'\right]^2 = \frac{m^2}{\left(1-c^2\right)}\sin^2\left(\frac{U}{2}\right) + K,\tag{5}$$

Where K is the constant of integration. Substituting  $K = 0, w(\xi) = \frac{U}{2}$  and  $a^2 = \frac{m^2}{(1-c^2)}$  in Eq. (5), it yields;

$$w' = a\sin(w). \tag{6}$$

Setting a = 1 in Eq. (6), gives;

$$w' = \sin(w). \tag{7}$$

Solving Eq. (7) by variables separable, we obtain the two significant equations as;

$$\sin(w) = \sin\left(w(\xi)\right) = \frac{2pe^{\xi}}{p^2 e^{2\xi} + 1}\Big|_{p=1} = \sec h(\xi),$$
(8)

$$\cos(w) = \cos\left(w(\xi)\right) = \frac{p^2 e^{2\xi} - 1}{p^2 e^{2\xi} + 1}\Big|_{p=1} = \tan h(\xi),$$
(9)

Where p is the integral constant and non-zero.

We suppose that the general nonlinear partial differential equation of the following form;

$$P(u, u_x, u_t, u_{xx}, u_{tt}, u_{xt}, ...) = 0, (10)$$

AIMS Mathematics

Volume 5, Issue 4, 4027–4044.

where u = u(x, t). We consider the solution of Eq. (10) as following expression,

$$U(\xi) = \sum_{i=1}^{n} \tanh^{i-1}(\xi) \Big[ B_i \sec h(\xi) + A_i \tanh(\xi) \Big] + A_0.$$
(11)

Eq. (11) can be rearranged according to Eq. (8) and Eq. (9) as follows;

$$U(w) = \sum_{i=1}^{n} \cos^{i-1}(w) \Big[ B_i \sin(w) + A_i \cos(w) \Big] + A_0.$$
(12)

Using the homogenous balance principle to determine the value of *n* between the highest power nonlinear term and highest derivative in the ordinary differential equation. We suppose that the summation of coefficients of  $\sin^i(w)\cos^j(w)$  with the same power is zero, this yields an equation system. With aid of the computational program, we solve the equation system to find the values of  $A_i, B_i, \mu$  and *c*. Finally, substituting the values of  $A_i, B_i, \mu$  and *c* into Eq. (11), we get the new travelling wave solutions to the Eq. (10).

#### 3. Applications of the method

In this section, we will present the application of the SGEM to the KP-BBM and the Benney–Luke equations.

#### 3.1. The KP-BBM equation

Let consider the following wave transform as,

$$u(x, y, t) = U(\zeta), \zeta = x + y - ct.$$
(13)

Using above transformation into Eq. (1), we get the following ordinary differential equation by integrating twice with respect to  $\zeta$ ,

$$(1+k-c)U - aU^{2} + bcU'' = 0.$$
(14)

We choose the integration constant to be zero. Balancing between U'' and  $U^2$  gives n = 2. For the value n = 2, the Eq. (12) takes the form,

$$U(w) = B_1 \sin(w) + A_1 \cos(w) + B_2 \cos(w) \sin(w) + A_2 \cos^2(w) + A_0.$$
 (15)

Differentiating Eq. (15) twice, yields

$$U''(w) = -2A_{1}\cos(w)\sin^{2}(w) - 4A_{2}\cos^{2}(w)\sin^{2}(w) + 2A_{2}\sin^{4}(w) + B_{1}\cos^{2}(w)\sin(w) - B_{1}\sin^{3}(w) + B_{2}\cos^{3}(w)\sin(w) - 5B_{2}\cos(w)\sin^{3}(w).$$
(16)

Substituting Eq. (15) and Eq. (16) into Eq. (14), we have a polynomial in term of trigonometric functions. Equating to zero all sum of coefficients of the same power of the trigonometric functions, we get the following algebraic equation system. Putting the values of the coefficients into Eq. (15), we obtain the following cases.

**Case 1:** When we consider as  $A_0 = -\frac{2b(1+k)}{a(1+b)}, A_2 = \frac{3b(1+k)}{a(1+b)}, B_2 = \frac{3ib(1+k)}{a(1+b)}, B_2 = \frac{3ib(1+k)}$ 

 $c = \frac{1+k}{1+b}$ ,  $A_1 = B_1 = 0$ , gives following complex mixed dark-bright soliton solution,

$$u_{1} = \frac{-2b(1+k)}{a(1+b)} - \frac{3ib(1+k)}{a(1+b)} Sech(f(x, y, t))Tanh(f(x, y, t)) + \frac{3b(1+k)}{a(1+b)}Tanh^{2}(f(x, y, t)), \quad (17)$$

where  $f(x, y, t) = \frac{(1+k)t}{1+b} - x - y$ . Considering suitable values of parameters, various figures may be observed as in Figures 1 and 2.



Figure 1. The complex mixed dark-bright soliton waves of Eq. (17) for the values of a = 0.6, b = 1, k = 0.5, y = 2.



**Figure 2.** The 2D-plots of Eq. (17) for the values of a = 0.6, b = 1, k = 0.5, y = 2, t = 0.1.

**Case 2:** When we take coefficients as,  $A_2 = -\frac{3A_0}{2}$ ,  $B_2 = \frac{3iA_0}{2}$ ,  $b = -\frac{aA_0}{2c}$ ,  $k = -1 + c - \frac{aA_0}{2}$ ,  $A_1 = B_1 = 0$ , gives the following,

$$u_{2} = A_{0} \left( 1 + \frac{-3e^{2ct} + 3e^{2(x+y)}}{2\left(e^{ct} - ie^{x+y}\right)^{2}} \right).$$
(18)

Considering suitable values of parameters, various figures may be observed as in Figures 3 and 4.



**Figure 3.** The complex combined dark-bright soliton waves of Eq. (18) for the values of c = 0.3,  $A_0 = 0.1$ , y = 1.2.

**AIMS Mathematics** 



**Figure 4.** The 2D-plots of Eq. (18) for the values of c = 0.3,  $A_0 = 0.1$ , y = 1.2, t = 0.11.

**Case 3:** If  $A_2 = -A_0, b = \frac{aA_0}{-6(1+k)+4aA_0}, c = 1+k-\frac{2aA_0}{3}, A_1 = B_1 = B_2 = 0$ , enable to write the solution as,

$$u_{3} = A_{0} - A_{0} \operatorname{Tanh}^{2} \left[ x + y - t \left( 1 + k - \frac{2aA_{0}}{3} \right) \right].$$
(19)

We consider suitable values of parameters at above solution, then some figures may be observed as in Figures 5 and 6.



**Figure 5.** The bright soliton waves of Eq. (19) for the values of  $A_0 = 3, a = 0.6, k = 0.5, y = 2$ .



**Figure 6.** The 2D-plot of Eq. (19) for the values of  $A_0 = 3, a = 0.6, k = 0.5, y = 2.$  t = 0.1.

AIMS Mathematics

Volume 5, Issue 4, 4027-4044.

**Case 4:** Considering as  $A_2 = -\frac{3A_0}{2}$ ,  $B_2 = \frac{3iA_0}{2}$ ,  $c = \frac{1+k}{1+b}$ ,  $a = \frac{-2b(1+k)}{A_0(1+b)}$ ,  $A_1 = B_1 = 0$ , we find the

$$u_{4} = -\frac{A_{0}\left(e^{\frac{2(1+k)t}{1+b}} - e^{2(x+y)} + 4ie^{\frac{2(1+k)t}{1+b} + x+y}\right)}{2\left(e^{\frac{2(1+k)t}{1+b}} - ie^{x+y}\right)^{2}}.$$
(20)

We found various figures for suitable values of parameters as Figures 7 and 8,



**Figure 7.** The complex solitary waves of Eq. (20) for the values of  $A_0 = 0.1, k = 0.3, b = -0.1, y = 1.2$ .



**Figure 8.** The 2D-plots of Eq. (20) for the values of  $A_0 = 0.1, k = 0.3, b = -0.1, y = 1.2, t = 0.1$ .

#### 3.2. The Benney-Luke equation

Let consider the following wave transform as,

$$u(x,t) = V(\zeta), \zeta = kx - ct.$$
(21)

Using above transformation into Eq. (2), we get the following nonlinear ordinary differential equation,

$$(c^{2}-k^{2})V'' + (\alpha k^{4} - \beta k^{2}c^{2})V^{4} - 3ck^{2}V'V'' = 0.$$
(22)

Integrating Eq. (22) with respect to  $\zeta$  once, we find,

$$2(c^{2}-k^{2})V'+2(\alpha k^{4}-\beta k^{2}c^{2})V'''-3ck^{2}(V')^{2}=0.$$
(23)

After getting transform V' = U, we can write as

$$2(c^{2}-k^{2})U+2(\alpha k^{4}-\beta k^{2}c^{2})U''-3ck^{2}U^{2}=0.$$
(24)

We choose the integration constant to be zero. Balancing between U'' and  $U^2$  gives n = 2.

Substituting Eq. (15) and Eq. (16) into Eq. (24), we have a polynomial in term of trigonometric functions.

Equating to zero all sum of coefficients of the same power of the trigonometric functions, we get the following algebraic equation system. Putting the values of the coefficients into Eq. (15), we obtain the following cases.

**Case 1:** When we consider 
$$A_0 = -\frac{2iB_2}{3}, A_2 = iB_2, \beta = -\frac{1}{2}i\left(\frac{B_2}{c} + \frac{4\alpha}{-2i+B_2c}\right), k = -\frac{(1+i)c}{\sqrt{2i-B_2c}}, A_1 = B_1 = 0,$$

gives the following complex mixed dark-bright soliton solution,

$$u_{1} = \frac{1}{3}iB_{2}\left(-ct - \frac{(1+i)cx}{\sqrt{2i - B_{2}c}} + 3i\,Sech\left[ct + \frac{(1+i)cx}{\sqrt{2i - B_{2}c}}\right] + 3Tanh\left[ct + \frac{(1+i)cx}{\sqrt{2i - B_{2}c}}\right]\right).$$
(25)

We consider suitable values of parameters to Eq. (25), then some figures may be observed as in Figures 9 and 10.



Figure 9. The complex combined dark-bright soliton waves of Eq. (25) for the values of  $c = 0.1, B_2 = 2$ .



**Figure 10.** The 2D-plots of Eq. (25) for the values of  $c = 0.1, B_2 = 2, t = 1$ .

**Case 2:** If we take following coefficients,  $A_0 = \frac{c^2 - k^2 + 5|c^2 - k^2|}{3ck^2}, A_2 = -\frac{2|c^2 - k^2|}{ck^2}, B_2 = \frac{2i}{c} - \frac{2ic}{k^2}, \alpha = \frac{-|c^2 - k^2| + c^2k^2\beta}{k^4}, A_1 = B_1 = 0,$  enable to write the solution as,

$$u_{2} = \frac{\left(c^{2} - k^{2} - \left|c^{2} - k^{2}\right|\right)\left(-ct + kx\right) + 6i\left(c^{2} - k^{2}\right)Sech\left[ct - kx\right] - 6\left|c^{2} - k^{2}\right|Tanh\left[ct - kx\right]}{3ck^{2}}.$$
 (26)

We consider suitable values of parameters at above solution, then some figures may be observed as in Figures 11 and 12.

**Figure 11.** The complex bright soliton waves of Eq. (26) for the values of c = 1, k = 0.5.



Figure 12: The 2D-plots of Eq. (26) for the values of c = 1, k = 0.5, t = 1.

**Case 3:** When the coefficients are taken as,  $A_0 = -\frac{2A_2}{3}$ ,  $B_2 = iA_2$ ,  $\alpha = \frac{(2+A_2c)(A_2+2c\beta)}{4c}$ ,  $k = -c\sqrt{\frac{2}{2+A_2c}}$ ,  $A_1 = B_1 = 0$ , we find another dark soliton solution,

$$u_{3} = \frac{1}{3}A_{2}\left(-ct - cx\sqrt{\frac{2}{2+A_{2}c}} - 3i\,Sech\left[ct + cx\sqrt{\frac{2}{2+A_{2}c}}\right] + 3Tanh\left[ct + cx\sqrt{\frac{2}{2+A_{2}c}}\right]\right). \tag{27}$$

AIMS Mathematics

Volume 5, Issue 4, 4027–4044.



Taking suitable values of parameters, various figures may be observed as in Figures 13 and 14.

**Figure 13.** The complex dark soliton waves of Eq. (27) for the values of c = 2,  $A_2 = 3$ .



**Figure 14.** The 2D-plots of Eq. (27) for the values of c = 2,  $A_2 = 3$ , t = 1.

Case 4: When we consider following coefficients,

$$A_{0} = \frac{4k(-\alpha + \beta)}{\sqrt{(-1 + 4k^{2}\alpha)(-1 + 4k^{2}\beta)}}, A_{2} = \frac{4k(\alpha - \beta)}{\sqrt{(-1 + 4k^{2}\alpha)(-1 + 4k^{2}\beta)}}, c = -k\sqrt{\frac{-1 + 4k^{2}\alpha}{-1 + 4k^{2}\beta}}, A_{1} = B_{1} = B_{2} = 0,$$

gives the following,

AIMS Mathematics

$$u_{4} = \frac{4k\left(-\alpha+\beta\right)Tanh\left[kx+kt\sqrt{\frac{-1+4k^{2}\alpha}{-1+4k^{2}\beta}}\right]}{\sqrt{\left(-1+4k^{2}\alpha\right)\left(-1+4k^{2}\beta\right)}}.$$
(28)

We found various figures for suitable values of parameters as above Figures 15 and 16.



**Figure 15.** The anti-kink wave soliton solution of Eq. (28) for the values of  $k = \beta = 1, \alpha = 2$ .



**Figure 16.** The 2D-plot of Eq. (28) for the values of  $k = \beta = 1, \alpha = 2, t = 1$ .

**Case** 5: 
$$A_0 = \frac{\sqrt{2}(\alpha - \beta)}{\sqrt{-\alpha - \beta}}, A_2 = \frac{\sqrt{2}(-\alpha + \beta)}{\sqrt{-\alpha - \beta}}, B_2 = \sqrt{\frac{2(\alpha - \beta)^2}{\alpha + \beta}}, c = \frac{\sqrt{-\alpha - \beta}}{\sqrt{2\beta}}, k = \frac{i}{\beta}, A_1 = B_1 = 0,$$

gives us,

$$u_{5} = \frac{\sqrt{2} \operatorname{Sech}\left[\frac{t\sqrt{-\alpha-\beta}}{\sqrt{2}\beta} - \frac{ix}{\sqrt{\beta}}\right] \left(-|\alpha-\beta|\sqrt{-\alpha-\beta} - (\alpha-\beta)\sqrt{\alpha+\beta} \operatorname{Sinh}\left[\frac{t\sqrt{-\alpha-\beta}}{\sqrt{2}\beta} - \frac{ix}{\sqrt{\beta}}\right]\right)}{i|\alpha+\beta|}.$$
(29)

Taking suitable values of parameters in Eq. (29), we found some simulations as in Figures17 and 18.

#### AIMS Mathematics



**Figure 17.** The complex dark waves of Eq. (29) for the values of  $\beta = -1, \alpha = 2$ .



**Figure 18.** The 2D-plots of Eq. (29) for the values of  $\beta = -1$ ,  $\alpha = 2$ , t = 1.

# 4. Results and discussion

We would like to thank you for following the instructions above very closely in advance. It will definitely save us lot of time and expedite the process of your paper's publication.

We consider the sine-Gordon expansion method which is based on sine-Gordon equation to construct new type solution to Eq. (1) and Eq. (2). In the literature, there have been many techniques

submitted such as Lie symmetry analysis, extended auxiliary equation mapping, extended direct algebraic mapping method, Hirota bilinear method, reductive perturbation technique. Kink, anti-kink solutions, lump solutions, line soliton, periodic waves and others have been shown by these mathematical tools. When we compare our results with other solutions, one can see that complex structure of obtained solutions are the feature that distinguishes it from others. Complex dark soliton, complex bright soliton solutions have more general structure than obtained by other techniques. In physics, complex calculus is considered to be of a formal mathematical, auxiliary type. Physical quantities can currently be defined by negative as well as by irrational number. Complex quantities extend some concepts as stability, convergence, conservation laws to reality. Consequently, complex-valued solutions can be verified in the experimental tests by physicist and engineers.

Complex numbers are particularly useful in explaining wave propagation. It is much easier to deal with the wave equation if we consider complex solutions. Since complex representation makes it much simpler to manage wave super-positions. If the refractive index of wave is enabled to be complex, the wave equation can be used to explain wave amplification and wave attenuation. In physical meaning, amplified waves are the fundamental basis for lasers when attenuated waves explain the absorption of light into black surfaces or the dissipation of sound into ambient heat [48].

Therefore, it is estimated that the results found in this paper may be used to explain such relationship of the governing models.

# 5. Conclusion

In this work, we have successfully performed SGEM to construct new complex combined dark-bright soliton solutions of the KP-BBM and the Benney–Luke equation. We have obtained many new soliton solutions. We observed that providing all solutions the mentioned two nonlinear partial differential equations by using a software. When we compare the SGEM with other analytical methods such as the extended tanh method, the new generalized (G'/G)-expansion method, the auxiliary equation method, we can deduce that this method is very effective and powerful tool for nonlinear models. To the best of our knowledge, these new complex combined dark-bright soliton solutions have been firstly submitted to the literature. It may be observed that wave behavior have shown their estimated wave propagations and distributions, physically, in Figures 1–18. Moreover, the submitted solutions may be helpful understanding physical phenomena especially in oceanography, geophysical science. The results show that the proposed method can be efficiently used to such completely integrable equations.

# **Conflict of interest**

The authors declare no conflict of interest.

### References

- 1. M. J. Ablowitz, C. W. Curtis, *Conservation laws and non-decaying for the Benney–Luke equation*, P. Roy. Soc. A Math. Phy., **469** (2013), 1–16.
- 2. D. Mitsotakis, *A simple introduction to water waves*, 2013, hal-00805080v1f. Available from: https://hal.archives-ouvertes.fr/hal-00805080v1.

- 3. A. M. Wazwaz, *The extended tanh method for new compact and noncompact solutions for the KP–BBM and the ZK–BBM equations*, Chaos Soliton. Fract., **38** (2008), 1505–1516.
- M. Song, C. Yang, B. Zhang, Exact solitary wave solutions of the Kadomtsov-Petviashvili-Benjamin-Bona-Mahony equation, Appl. Math. Comput., 217 (2010), 1334–1339.
- K. U. Tariq, A. R. Seadawy, Soliton solutions of (3+1)-dimensional Korteweg-de Vries Benjamin-Bona-Mahony, Kadomtsev-Petviashvili Benjamin-Bona-Mahony and modified Korteweg de Vries-Zakharov-Kuznetsov equations and their applications in water waves, J. King Saud Univ. Sci., 31 (2019), 8–13.
- 6. M. N. Alamand, M. A. Akbar, *Exact traveling wave solutions of the KP-BBM equation by using the new approach of generalized (G'/G)-expansion method*, SpringerPlus, **2** (2013), 1–7.
- 7. Z. Ouyang, *Traveling wave solutions of the Kadomtsev–Petviashvili–Benjamin–Bona–Mahony equation*, Abstr. Appl. Anal., **2014** (2014), 1–9.
- J. Q. Lü, S. Bilige, X. Q. Gao, et al. Abundant lump solutions and interaction phenomena to the Kadomtsev-Petviashvili-Benjamin-Bona-Mahony equation, J. Appl. Math. Phys., 6 (2018), 1733–1747.
- 9. J. Akter, M. A. Akbar, *Exact solutions to the Benney–Luke equation and the Phi-4 equations by using modified simple equation method*, Results Phys., **5** (2015), 125–130.
- 10. S. Sirisubtawee, S. Koonprasert, *Exact traveling wave solutions of certain nonlinear partial differential equations using the*  $(G'/G^2)$ *-expansion method*, Adv. Math. Phys., **2018** (2018), 1–16.
- 11. D. J. Benney, J. C. Luke, *On the interactions of permanent waves of finite amplitude*, J. Math. Phys., **43** (1964), 309–313.
- 12. H. O. Roshid, M. A. Rahman, *The exp*  $(-\varphi(\eta))$ *-expansion method with application in the* (1+1)*- dimensional classical Boussinesq equations*, Results Phys., **4** (2014), 150–155.
- 13. S. Wang, G. Xu, G. Chen, *Cauchy problem for the generalized Benney–Luke equation*, J. Math. Phys., **48** (2007), 073521.
- 14. J. R. Quintero, J. C. M. Grajales, *Instability of solitary waves for a generalized Benney–Luke equation*, Nonlinear Anal. Theor., **68** (2008), 3009–3033.
- 15. H. Triki, A. Yildirim, T. Hayat, et al. *Shock wave solution of the Benney-Luke equation*, Romanian J. Phys., **57** (2012), 1029–1034.
- 16. R. Hirota, The Direct Method in Soliton Theory, Cambridge University Press, Cambridge, 2004.
- 17. M. J. Ablowitz, P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering*, Cambridge University Press, Cambridge, 1991.
- H. Triki, T. Hayat, O. M. Aldossary, et al. *1-Soliton solution of the three component system of Wu-Zhang equations*, Hacet. J. Math. Stat., 41 (2012), 537–543.
- 19. A. M. Wazwaz, H. Triki, Soliton solutions for a generalized KdV and BBM equations with time-dependent coefficients, Commun. Nonlinear Sci., 16 (2011), 1122–1126.
- 20. W. Malfliet, W. Hereman, *The tanh method: II. Perturbation technique for conservative systems*, Phys. Scr., **54** (1996), 569.
- M. Iqbal, A. R. Seadawy, D. Lu, Construction of solitary wave solutions to the nonlinear modified Kortewege-de Vries dynamical equation in unmagnetized plasma via mathematical methods, Mod. Phys. Lett. A, 33 (2018), 1–13.

- 22. D. Lu, A. R. Seadawy, M. Iqbal, *Mathematical methods via construction of traveling and solitary wave solutions of three coupled system of nonlinear partial differential equations and their applications*, Results Phys., **11** (2018), 1161–1171.
- M. Iqbal, A. R. Seadawy, D. Lu, Dispersive solitary wave solutions of nonlinear further modified Korteweg-de Vries dynamical equation in an unmagnetized dusty plasma, Mod. Phys. Lett. A, 33 (2018), 1–19.
- 24. A. R. Seadawy, D. Lu, M. Iqbal, Application of mathematical methods on the system of dynamical equations for the ion sound and Langmuir waves, Pramana J. Phys., **93** (2019), 10.
- M. Iqbal, A. R. Seadawy, D. Lu, Applications of nonlinear longitudinal wave equation in a magneto-electro-elastic circular rod and new solitary wave solutions, Mod. Phys. Lett. B, 33 (2019), 1–17.
- A. R. Seadawy, M. Iqbal, D. Lu, Applications of propagation of long-wave with dissipation and dispersion in nonlinear media via solitary wave solutions of generalized Kadomtsev–Petviashvili modified equal width dynamical equation, Comput. Math. Appl., 78 (2019), 3620–3632.
- 27. M. Iqbal, A. R. Seadawy, D. Lu, et al. *Construction of a weakly nonlinear dispersion solitary* wave solution for the Zakharov-Kuznetsov-modified equal width dynamical equation, Indian J. Phys., 2019, 1–10.
- M. Iqbal, A. R. Seadawy, D. Lu, et al. Construction of bright-dark solitons and ion-acoustic solitary wave solutions of dynamical system of nonlinear wave propagation, Mod. Phys. Lett. A, 34 (2019), 1–24.
- 29. A. R. Seadawy, D. Lu, N. Nasreen, Construction of solitary wave solutions of some nonlinear dynamical system arising in nonlinear water wave models, Indian J. Phys., 2019, 1–10.
- 30. A. R. Seadawy, M. Iqbal, D. Lu, Nonlinear wave solutions of the Kudryashov–Sinelshchikov dynamical equation in mixtures liquid-gas bubbles under the consideration of heat transfer and viscosity, J. Taibah Univ. Sci., **13** (2019), 1060–1072.
- 31. A. R. Seadawy, M. Iqbal, D. Lu, Propagation of kink and anti-kink wave solitons for the nonlinear damped modified Korteweg-de Vries equation arising in ion-acoustic wave in an unmagnetized collisional dusty plasma, Physica A, **544** (2019), 123560.
- 32. C. Q. Dai, Y. Fan, N. Zhang, *Re-observation on localized waves constructed by variable separation solutions of (1+1)-dimensional coupled integrable dispersionless equations via the projective Riccati equation method*, Appl. Math. Lett., **96** (2019), 20–26.
- H. An, D. Feng, H. Zhu, General, M-lump, high-order breather and localized interaction solutions to the (2+1)-dimensional Sawada–Kotera equation, Nonlinear Dyn., 98 (2019), 1275–1286.
- J. J. Yang, S. F. Tian, W. Q. Peng, et al. *The N-coupled higher-order nonlinear Schrödinger equation: Riemann-Hilbert problem and multi-soliton solutions*, Math. Meth. Appl. Sci., 43 (2020), 2458–2472.
- W. Q. Peng, S. F. Tian, X. B. Wang, et al. Riemann-Hilbert method and multi-soliton solutions for three-component coupled nonlinear Schrödinger equations, J. Geom. Phys., 146 (2019), 1–9.
- 36. T. T. Zhang, On Lie symmetry analysis, conservation laws and solitary waves to a longitudinal wave motion equation, Appl. Math. Lett., **98** (2019), 199–205.

- 37. S. F. Tian, Lie symmetry analysis, conservation laws and solitary wave solutions to a fourth-order nonlinear generalized Boussinesq water wave equation, Appl. Math. Lett., 100 (2020), 106056.
- E. İ. Eskitaşçıoğlu, M. B. Aktaş, H. M. Baskonus, New complex and hyperbolic forms for Ablowitz-Kaup-Newell-Segur wave equation with fourth order, Appl. Math. Nonlinear Sci., 4 (2019), 105–112.
- 39. W. Gao, B. Ghanbari, H. Günerhan, et al. Some mixed trigonometric complex soliton solutions to the perturbed nonlinear Schrödinger equation, Mod. Phys. Lett. B, **34** (2020), 1–18.
- 40. W. Gao, H. F. Ismael, H. Bulut, et al. *Instability modulation for the* (2+1)-dimension paraxial wave equation and its new optical soliton solutions in Kerr media, Phys. Scr., **95** (2020), 1–12.
- 41. A. Cordero, J. P. Jaiswal, J. R. Torregrosa, *Stability analysis of fourth-order iterative method for finding multiple roots of non-linear equations*, Appl. Math. Nonlinear Sci., **4** (2019), 43–56.
- L. D. Moleleki, T. Motsepa, C. M. Khalique, Solutions and conservation laws of a generalized second extended (3+1)-dimensional Jimbo–Miwa equation, Appl. Math. Nonlinear Sci., 3 (2018), 459–474.
- 43. P. K. Pandey, A new computational algorithm for the solution of second order initial value problems in ordinary differential equations, Appl. Math. Nonlinear Sci., **3** (2018), 167–174.
- 44. C. Yan, A simple transformation for nonlinear waves, Phys. Lett. A, 224 (1996), 77-84.
- 45. Z. Yan, H. Zhang, New explicit and exact travelling wave solutions for a system of variant Boussinesq equations in mathematical physics, Phys. Lett. A, **252** (1999), 291–296.
- 46. G. Yel, H. M. Baskonus, H. Bulut, Novel archetypes of new coupled Konno–Oono equation by using sine–Gordon expansion method, Opt. Quant. Electron., **49** (2017), 285.
- 47. H. M. Baskonus, H. Bulut, T. A. Sulaiman, *Investigation of various travelling wave solutions to the extended* (2+1)-*dimensional quantum ZK equation*, Eur. Phys. J. Plus, **132** (2017), 1–10.
- 48. Y. D. Chong, *MH2801: Complex Methods for the Sciences*, Nanyang Technological University, 2016. Available from: http://www1.spms.ntu.edu.sg/~ydchong/teaching.html.



© 2020 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0)