



Research article

Mild solutions for a multi-term fractional differential equation via resolvent operators

Yong-Kui Chang¹ and Rodrigo Ponce^{2,*}

¹ School of Mathematics and Statistics, Xidian University, Xi'an 710071, Shaanxi, P. R. China

² Universidad de Talca, Instituto de Matemáticas, Casilla 747, Talca-Chile

* **Correspondence:** Email: rponce@inst-mat.otalca.cl.

Abstract: This paper is concerned with multi-term fractional differential equations. With the help of the theory of fractional resolvent families, we establish the existence of mild solutions to a multi-term fractional differential equation.

Keywords: Caputo fractional derivatives; nonlocal problem; fractional diffusion equation; resolvent operators

Mathematics Subject Classification: 33E12, 34A08, 45N05, 47D06, 34K30

1. Introduction

In the last two decades, differential equations involving fractional derivatives, have been used in many mathematical models to describe a wide variety of phenomena, including problems in viscoelasticity, signal and image processing, engineering, economics, epidemiology and among others, and the study of this kind of equations has been a topic of interest in recent years. See [9, 16, 19, 25, 37, 41–43, 45] and the references therein.

In this paper, we consider the following multi-term fractional differential equations

$$\partial^\alpha u(t) = Au(t) + \partial^{\alpha-\beta} f(t, u(t)), \quad t \in \mathbb{R}, \tag{1.1}$$

and

$$\partial_t^\alpha u(t) = Au(t) + \partial_t^{\alpha-\beta} f(t, u(t)), \quad t \in [0, T], \tag{1.2}$$

where A is a closed linear operator defined in a Banach space X , $1 < \alpha, \beta < 2$, $T > 0$, and f is a suitable continuous function. Here, for $\gamma > 0$ the derivatives $\partial^\gamma u$ and $\partial_t^\gamma u$, denote the Weyl and Caputo fractional derivatives, respectively.

Although the definition of the fractional derivatives in the sense of Weyl (defined on \mathbb{R}) and Caputo (defined on $[0, \infty)$) are different, we notice that the mild solution to Eqs (1.1) and (1.2) can be written in terms of the same resolvent family. In fact, if A is the generator of the fractional resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$ (see its definition in Section 2) then the *mild* solutions to Eqs (1.1) and (1.2) are defined, respectively, by

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in \mathbb{R},$$

and

$$u(t) = S_{\alpha,1}(t)x + S_{\alpha,2}(t)y + \int_0^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t > 0,$$

where $x = u(0)$ and $y = u'(0)$ are the initial conditions in Eq (1.2), and the families $\{S_{\alpha,\beta}(t)\}_{t \geq 0}$, and $\{S_{\alpha,2}(t)\}_{t \geq 0}$, are given respectively by

$$S_{\alpha,\beta}(t) = (g_{\beta-1} * S_{\alpha,1})(t), \quad \text{and} \quad S_{\alpha,2}(t) = (g_1 * S_{\alpha,1})(t).$$

Here, the $*$ denotes the usual finite convolution and for $\gamma > 0$ the function g_γ is defined by $g_\gamma(t) := t^{\gamma-1}/\Gamma(\gamma)$, where $\Gamma(\cdot)$ is the Gamma function. The fractional resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$ is defined by

$$S_{\alpha,1}(t) := \frac{1}{2\pi i} \int_{\Gamma} e^{\lambda t} \lambda^{\alpha-1} (\lambda^\alpha - A)^{-1} d\lambda, \quad t \geq 0,$$

where Γ is a suitable complex path where the resolvent operator $(\lambda^\alpha - A)^{-1}$ is well defined. By the uniqueness of the Laplace transform it is easy to see that

$$S_{\alpha,2}(t) = \frac{1}{2\pi i} \int_{\Gamma} e^{\lambda t} \lambda^{\alpha-2} (\lambda^\alpha - A)^{-1} d\lambda \quad \text{and} \quad S_{\alpha,\beta}(t) = \frac{1}{2\pi i} \int_{\Gamma} e^{\lambda t} \lambda^{\alpha-\beta} (\lambda^\alpha - A)^{-1} d\lambda,$$

for all $t \geq 0$. The existence of mild solutions to Eq (1.1) in case $\beta = 1$ has been widely studied in the last years, see for instance [4, 12, 13, 24] and references therein. In these mentioned papers, the operator A is assumed to be an ω -sectorial operator of angle θ (see definition in Section 2). In this case, A generates a resolvent family $\{E_\alpha(t)\}_{t \geq 0}$ (see [11, 28]) which satisfies

$$\|E_\alpha(t)\| \leq \frac{C}{1 + |\omega|t^\alpha}, \quad \text{for all } t \geq 0,$$

where C is a positive constant depending only on α and θ . This decay of $\{E_\alpha(t)\}_{t \geq 0}$ provides also some tools to obtain many and interesting consequences on the study of qualitative properties of solutions to fractional (and integral) differential (and difference) equations. See for instance [4, 7, 8, 29, 31, 44] and the references therein for further details. We notice that, by the uniqueness of the Laplace transform, the resolvent families $\{E_\alpha(t)\}_{t \geq 0}$ and $\{S_{\alpha,1}(t)\}_{t \geq 0}$ are the same for $1 < \alpha < 2$.

On the other hand, the existence of mild solutions to fractional differential equations with nonlocal conditions has been studied by several authors in the last years. The concept of nonlocal initial condition was introduced by L. Byszewski [6] to extend the study of classical initial value problems. This notion results more suitable to describe more precisely several phenomena in applied sciences, because it considers additional information in the initial data. More concretely, the nonlocal conditions have the form $u(0) + g(u) = u_0$ instead $u(0) = u_0$, where g is an appropriate function that

represents the additional information in the system and provides a better description of the initial state of the system than the classical initial value problem. The theory of nonlocal Cauchy problems has been developed rapidly and has been studied widely in the last years, see for instance [3, 33, 38] and the references therein for more details.

There exists a wide recent literature on the existence of mild solutions to fractional differential equations with nonlocal initial conditions. More specifically, the problem

$$\begin{cases} \partial_t^\alpha u(t) &= Au(t) + f(t, u(t)), \quad t \in [0, T] \\ u(0) + g(u) &= u_0, \end{cases} \quad (1.3)$$

where $T > 0$, A is a closed linear operator defined in a Banach space X , $0 < \alpha \leq 1$, $u_0 \in X$, f is a suitable semilinear continuous function has been studied extensively in recent years. See for instance [1, 2, 10, 26, 30, 35]. Since the fractional derivative ∂_t^α for $\alpha = 1$ is the usual derivative $\frac{d}{dt}$, the case $\alpha = 1$ in (1.3) corresponds precisely to the semilinear Cauchy problem introduced in the seminal paper [6] and the theory of C_0 -semigroups of linear operators is the main tool to obtain the existence of solutions in this case. Similarly, for $\alpha > 0$ the theory of fractional resolvent families represents one of the main tools to study the existence of mild solutions to (1.3). Indeed, if $0 < \alpha \leq 1$ and A generates a resolvent family $\{S_{\alpha, \alpha}(t)\}_{t \geq 0}$, then the mild solution to (1.3) is given by

$$u(t) = S_{\alpha, 1}(u_0 - g(u)) + \int_0^t S_{\alpha, \alpha}(t-s)f(s, u(s))ds \quad (1.4)$$

where $S_{\alpha, 1}(t) := (g_{1-\alpha} * S_{\alpha, \alpha})(t)$, see for instance [30]. We notice that the variation of constant formula (1.4) coincides with the case $\alpha = 1$ introduced in [6, Section 3]. Similarly, for $1 < \alpha < 2$ and $\beta = 1$ or $\beta = \alpha$, the Eq (1.2) subject to the nonlocal conditions $u(0) + g(u) = u_0$, and $u'(0) + h(u) = u_1$, where $g, h : C(I, X) \rightarrow X$ are continuous and u_0, u_1 belong to X , ($I := [0, T]$) has been considered by several authors in the last years. See for instance [2, 22] for the case $\beta = 1$ and [33, 34] in case $\beta = \alpha$.

In this paper, our concern is the study of existence of mild solutions to the fractional differential Eqs (1.1) and (1.2). Here, we assume certain conditions on the operator A and on the parameters α and β in order to ensure that A is the generator of a fractional resolvent family $\{S_{\alpha, \beta}(t)\}_{t \geq 0}$.

More specifically, in Eq (1.1) we consider the Weyl fractional derivative, because it is defined for functions on \mathbb{R} . More precisely, we show that if the function f in (1.1) is an almost periodic or an almost automorphic (among others) vector-valued function, then the Eq (1.1) has a unique almost periodic or almost automorphic function mild solution, respectively, which is given in terms of $\{S_{\alpha, \beta}(t)\}_{t \geq 0}$.

On the other hand, in Eq (1.2) the derivative is taken in the sense of Caputo, because it is defined on the positive real axis $[0, \infty)$. Under the the nonlocal conditions $u(0) + g(u) = u_0$, and $u'(0) + h(u) = u_1$ we prove that (1.2) has at least one mild solution. Here, the properties of the fractional resolvent family $\{S_{\alpha, \beta}(t)\}_{t \geq 0}$ are again an important tool to obtain the result.

This paper is organized as follows. The Section 2 gives the preliminaries on fractional calculus, sectorial operators, fractional resolvent families and some subspaces of bounded and continuous functions. Section 3 is devoted to the existence of mild solutions to (1.1). In Section 4 is studied the existence of mild solutions to the nonlocal problem (1.2). Finally, in Section 5 we give some examples.

2. Preliminaries

For a Banach space $(X, \|\cdot\|)$, the space of all bounded and linear operators from X into X is denoted by $\mathcal{B}(X)$. If A is a closed linear operator defined on X we denote by $\rho(A)$ the resolvent set of A and $R(\lambda, A) = (\lambda - A)^{-1}$ its resolvent operator, which is defined for all $\lambda \in \rho(A)$. For $1 \leq p < \infty$, $L^p(\mathbb{R}_+, X)$ denotes the space of all Bochner measurable functions $g : \mathbb{R}_+ \rightarrow X$ such that

$$\|g\|_p := \left(\int_0^\infty \|g(t)\|^p dt \right)^{1/p} < \infty.$$

We recall that a strongly continuous family $\{S(t)\}_{t \geq 0} \subset \mathcal{B}(X)$ is said to be exponentially bounded if there exist two constants $M > 0$ and $w \in \mathbb{R}$ such that $\|S(t)\| \leq Me^{wt}$ for all $t > 0$.

A closed and densely defined operator A , defined on a Banach space $(X, \|\cdot\|)$, is said to be ω -sectorial of angle ϕ , if there exist $\phi \in [0, \pi/2)$ and $\omega \in \mathbb{R}$ such that its resolvent exists in the sector $\omega + \Sigma_\phi := \{\omega + \lambda : \lambda \in \mathbb{C}, |\arg(\lambda)| < \frac{\pi}{2} + \phi\} \setminus \{\omega\}$ and $\|R(\lambda, A)\| \leq \frac{M}{|\lambda - \omega|}$ for all $\lambda \in \omega + \Sigma_\phi$. See [17] and [18] for further details.

Now, we review some results on fractional calculus. We recall that for $\gamma > 0$, the function g_γ is defined by $g_\gamma(t) = \frac{t^{\gamma-1}}{\Gamma(\gamma)}$ for all $t \geq 0$. For $\gamma > 0$, $[\gamma]$ denotes the smallest integer greater than or equal to γ , and $[\gamma]$ denotes the integer part of γ . As usual, the finite convolution of f and g is defined by $(f * g)(t) = \int_0^t f(t-s)g(s)ds$.

Definition 1. Let $\alpha > 0$ and $n = [\alpha]$. The Caputo fractional derivative of order α of a function $u : [0, \infty) \rightarrow X$ is defined by

$$\partial_t^\alpha u(t) := \int_0^t g_{n-\alpha}(t-s)u^{(n)}(s)ds.$$

Definition 2. Let $\alpha > 0$ and $n = [\alpha] + 1$. The Weyl fractional derivative of order α of a function $u : \mathbb{R} \rightarrow X$ is defined by

$$\partial_t^\alpha u(t) := \frac{d^n}{dt^n} \partial^{-(n-\alpha)} u(t),$$

where for $\gamma > 0$, $\partial^{-\gamma} u(t) := \int_{-\infty}^t g_\gamma(t-s)u(s)ds$ for all $t \in \mathbb{R}$.

It is a well known fact that if $\alpha \in \mathbb{N}$, then $\partial_t^\alpha = \partial^n = \frac{d^n}{dt^n}$, that is, the Caputo and Weyl fractional derivatives coincide with the usual derivative if $\alpha \in \mathbb{N}$. Moreover, if $\alpha, \beta \in \mathbb{R}$, then $\partial^\alpha \partial^\beta u = \partial^\beta \partial^\alpha u = \partial^{\alpha+\beta} u$. See [25, 41] and [42] for more details and applications on fractional differential calculus.

Now, we recall the resolvent families of operators generated by an operator A .

Definition 3. Let A be closed linear operator with domain $D(A)$, defined on a Banach space X , $1 \leq \alpha \leq 2$ and $0 < \beta \leq 2$. We say that A is the generator of an (α, β) -resolvent family, if there exists $\nu \geq 0$ and a strongly continuous and exponentially bounded function $S_{\alpha, \beta} : [0, \infty) \rightarrow \mathcal{B}(X)$ such that $\{\lambda^\alpha : \operatorname{Re} \lambda > \nu\} \subset \rho(A)$, and for all $x \in X$,

$$\lambda^{\alpha-\beta} (\lambda^\alpha - A)^{-1} x = \int_0^\infty e^{-\lambda t} S_{\alpha, \beta}(t) x dt, \quad \operatorname{Re} \lambda > \nu.$$

In this case, $\{S_{\alpha, \beta}(t)\}_{t \geq 0}$ is called the (α, β) -resolvent family generated by A .

If we compare Definition 3 with the notion of (a, k) -regularized families introduced in [21], then we notice that $t \mapsto S_{\alpha, \beta}(t)$, is a (g_α, g_β) -regularized family. Moreover, the family $\{S_{\alpha, \beta}(t)\}_{t \geq 0}$ is well known in some cases. For example, $S_{1,1}(t)$ is a C_0 -semigroup, $S_{2,1}(t)$, corresponds to a cosine family and $S_{2,2}(t)$ is a sine family. In the scalar case, that is, when $A = \varrho \mathcal{I}$, where $\varrho \in \mathbb{C}$ and \mathcal{I} denotes the identity operator, then by the uniqueness of the Laplace transform, $S_{\alpha, \beta}(t)$ corresponds to the function $t^{\beta-1} E_{\alpha, \beta}(\varrho t^\alpha)$, where for $z \in \mathbb{C}$ the generalized Mittag-Leffler function is defined by $E_{\alpha, \beta}(z) = \sum_{k=0}^{\infty} \frac{z^k}{\Gamma(\alpha k + \beta)}$, see for instance [39, 40]. See also [41] and [42] for an interesting and recent discussion on the theory of general fractional derivatives and its applications.

We have also the following result. Its proof follows similarly as in [20, Proposition 3.7].

Proposition 4. *Let $1 \leq \alpha, \beta \leq 2$. Let $S_{\alpha, \beta}(t)$ be the (α, β) -resolvent family generated by A . Then:*

1. $S_{\alpha, \beta}(t)x \in D(A)$ and $S_{\alpha, \beta}(t)Ax = AS_{\alpha, \beta}(t)x$ for all $x \in D(A)$ and $t \geq 0$.
2. If $x \in D(A)$ and $t \geq 0$, then

$$S_{\alpha, \beta}(t)x = g_\beta(t)x + \int_0^t g_\alpha(t-s)AS_{\alpha, \beta}(s)x ds \quad (2.1)$$

3. If $x \in X, t \geq 0$, then $\int_0^t g_\alpha(t-s)S_{\alpha, \beta}(s)x ds \in D(A)$ and $S_{\alpha, \beta}(t)x = g_\beta(t)x + A \int_0^t g_\alpha(t-s)S_{\alpha, \beta}(s)x ds$.

In particular, $S_{\alpha, \beta}(0) = g_\beta(0)\mathcal{I}$.

The next result gives sufficient conditions on α, β and A to obtain generators of (α, β) -resolvent families.

Theorem 5. [28] *Let $1 < \alpha < 2$ and $\beta \geq 1$ such that $\alpha - \beta + 1 > 0$. Assume that A is ω -sectorial of angle $\frac{(\alpha-1)\pi}{2}$, where $\omega < 0$. Then A generates an exponentially bounded (α, β) -resolvent family.*

Theorem 6. [28] *Let $1 < \alpha < 2$ and $\beta \geq 1$ such that $\alpha - \beta + 1 > 0$. Assume that A is ω -sectorial of angle $\frac{(\alpha-1)\pi}{2}$, where $\omega < 0$. Then, there exists a constant $C > 0$, depending only on α and β , such that*

$$\|S_{\alpha, \beta}(t)\| \leq \frac{Ct^{\beta-1}}{1 + |\omega|t^\alpha}, \quad \text{for all } t > 0. \quad (2.2)$$

Finally, we recall some spaces of functions. For a given Banach space $(X, \|\cdot\|)$, let $BC(X) := \{f : \mathbb{R} \rightarrow X : \|f\|_\infty := \sup_{t \in \mathbb{R}} \|f(t)\| < \infty\}$ be the Banach space of all bounded and continuous functions. For $T > 0$ fixed, $P_T(X)$ denotes the space of all vector-valued periodic functions, that is, $P_T(X) := \{f \in BC(X) : f(t+T) = f(t), \text{ for all } t \in \mathbb{R}\}$. We denote by $AP(X)$ to the space of all almost periodic functions (in the sense of Bohr), which consists of all $f \in BC(X)$ such that for every $\varepsilon > 0$ there exists $l > 0$ such that for every subinterval of \mathbb{R} of length l contains at least one point τ such that $\|f(t+\tau) - f(t)\|_\infty \leq \varepsilon$. A function $f \in BC(X)$ is said to be almost automorphic if for every sequence of real numbers $(s'_n)_{n \in \mathbb{N}}$ there exists a subsequence $(s_n)_{n \in \mathbb{N}} \subset (s'_n)_{n \in \mathbb{N}}$ such that

$$g(t) := \lim_{n \rightarrow \infty} f(t + s_n)$$

is well defined for each $t \in \mathbb{R}$, and

$$f(t) = \lim_{n \rightarrow \infty} g(t - s_n), \quad \text{for each } t \in \mathbb{R}.$$

We denote by $AA(X)$ the Banach space of all almost automorphic functions.

On the other hand, the space of compact almost automorphic functions is the space of all functions $f \in BC(X)$ such that for all sequence $(s'_n)_{n \in \mathbb{N}}$ of real numbers there exists a subsequence $(s_n)_{n \in \mathbb{N}} \subset (s'_n)_{n \in \mathbb{N}}$ such that $g(t) := \lim_{n \rightarrow \infty} f(t + s_n)$ and $f(t) = \lim_{n \rightarrow \infty} g(t - s_n)$ uniformly over compact subsets of \mathbb{R} .

We notice that $P_T(X), AP(X), AA(X)$ and $AA_c(X)$ are Banach spaces under the norm $\|\cdot\|_\infty$ and

$$P_T(X) \subset AP(X) \subset AA(X) \subset AA_c(X) \subset BC(X).$$

We notice that all these inclusions are proper. Now we consider the set $C_0(X) := \{f \in BC(X) : \lim_{|t| \rightarrow \infty} \|f(t)\| = 0\}$, and define the space of asymptotically periodic functions as $AP_T(X) := P_T(X) \oplus C_0(X)$. Analogously, we define the space of asymptotically almost periodic functions,

$$AAP(X) := AP(X) \oplus C_0(X),$$

the space of asymptotically compact almost automorphic functions,

$$AAA_c(X) := AA_c(X) \oplus C_0(X),$$

and the space of asymptotically almost automorphic functions,

$$AAA(X) := AA(X) \oplus C_0(X).$$

We have the following natural proper inclusions

$$AP_T(X) \subset AAP(X) \subset AAA_c(X) \subset AAA(X) \subset BC(X).$$

For more details on this function spaces, we refer to reader to [23, 27].

Throughout, we will use the notation $\mathcal{N}(X)$ to denote any of the function spaces $AP_T(X), AAP(X), AAA_c(X)$ and $AAA(X)$ defined above. Finally, we define the set $\mathcal{N}(\mathbb{R} \times X; X)$ which consists of all functions $f : \mathbb{R} \times X \rightarrow X$ such that $f(\cdot, x) \in \mathcal{N}(X)$ uniformly for each $x \in K$, where K is any bounded subset of X . Moreover, we have the following result.

Theorem 7. [23] Let $\{S(t)\}_{t \geq 0} \subset \mathcal{B}(X)$ be a strongly continuous and uniformly 1-integrable family, that is $\int_0^\infty \|S(t)\| dt < \infty$. If $f \in \mathcal{N}(X)$, then the function $u : \mathbb{R} \rightarrow X$ defined by

$$u(t) := \int_{-\infty}^t S(t-s)f(s)ds,$$

belongs to $\mathcal{N}(X)$.

3. Bounded mild solutions to Eq (1.1)

Let $1 < \alpha < 2$ and $\beta \geq 1$. In this section, we first consider the linear version of the Eq (1.1), that is,

$$\partial^\alpha u(t) = Au(t) + \partial^{\alpha-\beta} f(t), \quad t \in \mathbb{R}. \quad (3.1)$$

Definition 8. A function $u \in C(\mathbb{R}, X)$ is called a mild solution to Eq (3.1) if the function $s \mapsto S_{\alpha,\beta}(t-s)f(s)$ is integrable on $(-\infty, t)$ for each $t \in \mathbb{R}$ and

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds, \quad t \in \mathbb{R}. \quad (3.2)$$

We notice that (3.1) can be considered as the limiting equation of the following integro-differential equation with singular kernels

$$\begin{cases} v'(t) = \int_0^t \frac{(t-s)^{\alpha-2}}{\Gamma(\alpha-1)} Av(s) + \frac{(t-s)^{\beta-2}}{\Gamma(\beta-1)} f(s)ds, & t \geq 0 \\ v(0) = v_0, & v_0 \in X, \end{cases} \quad (3.3)$$

in the sense that the mild solution to Eq (3.3) converges to the mild solution of (3.1) as $t \rightarrow \infty$. In fact, if $\omega < 0$ and A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)\pi}{2}$, then taking Laplace transform in (3.3) we obtain

$$\lambda \hat{v}(\lambda) - v(0) = \frac{1}{\lambda^{\alpha-1}} A \hat{v}(\lambda) + \frac{1}{\lambda^{\beta-1}} \hat{f}(\lambda), \quad \operatorname{Re} \lambda > 0,$$

which is equivalent to

$$(\lambda^\alpha - A) \hat{v}(\lambda) = \lambda^{\alpha-1} v(0) + \lambda^{\alpha-\beta} \hat{f}(\lambda), \quad \operatorname{Re} \lambda > 0.$$

Therefore the solution of problem (3.3) can be written as

$$v(t) = S_{\alpha,1}(t)v_0 + \int_0^t S_{\alpha,\beta}(t-s)f(s)ds, \quad t \geq 0, \quad (3.4)$$

where $\{S_{\alpha,\beta}(t)\}_{t \geq 0}$ is the family of operators given by

$$S_{\alpha,\beta}(t) := (g_{\beta-1} * S_{\alpha,1})(t).$$

On the other hand, by [28, Corollary 3.9] the function $t \mapsto S_{\alpha,\beta}(t)$ is uniformly 1-integrable and therefore if f is a bounded continuous function (for example, if f belongs to $\mathcal{N}(X)$), then the mild solution to Eq (1.1) is given by

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds.$$

Since

$$v(t) - u(t) = S_{\alpha,1}(t)v_0 - \int_t^\infty S_{\alpha,\beta}(s)f(t-s)ds,$$

we conclude by [28, Corollary 3.8], that $v(t) - u(t) \rightarrow 0$ as $t \rightarrow \infty$.

Let $1 < \alpha < 2$, $\beta \geq 1$ such that $\alpha - \beta + 1 > 0$, $\omega < 0$ and assume that A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)\pi}{2}$. By Theorem 5, the operator A generates a resolvent family $\{S_{\alpha,\beta}(t)\}_{t \geq 0}$. Take a bounded and continuous function $f : \mathbb{R} \rightarrow X$, (for example, we can take $f \in \mathcal{N}(X)$). Define the function $\phi(t)$ by

$$\phi(t) := \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds, \quad t \in \mathbb{R}. \quad (3.5)$$

By Theorem 6 we have $\|\phi\|_\infty \leq \|S_{\alpha,\beta}\|_1 \|f\|_\infty$. If $f(t) \in D(A)$ for all $t \in \mathbb{R}$, then $\phi(t) \in D(A)$ for all $t \in \mathbb{R}$ (see [5, Proposition 1.1.7]). Assume that $\partial^\alpha \phi$ exists. The Proposition 4 and Fubini's theorem imply that

$$\begin{aligned}
 \partial^\alpha \phi(t) &= \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s)\phi(s)ds \\
 &= \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s S_{\alpha,\beta}(s-r)f(r)drds \\
 &= \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s \left[g_\beta(s-r)f(r) + (g_\alpha * AS_{\alpha,\beta})(s-r)f(r) \right] drds \\
 &= \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s)\partial^{-\beta}f(s)ds + \\
 &\quad \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s \int_0^{s-r} g_\alpha(s-r-v)AS_{\alpha,\beta}(v)f(r)dvdrds \\
 &= \partial^{\alpha-\beta}f(t) + \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s \int_r^s g_\alpha(s-w)AS_{\alpha,\beta}(w-r)f(r)dwdrds \\
 &= \partial^{\alpha-\beta}f(t) + \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s \int_{-\infty}^w g_\alpha(s-w)AS_{\alpha,\beta}(w-r)f(r)drdwds \\
 &= \partial^{\alpha-\beta}f(t) + \frac{d^n}{dt^n} \int_{-\infty}^t g_{n-\alpha}(t-s) \int_{-\infty}^s g_\alpha(s-w)A\phi(w)dwds \\
 &= \partial^{\alpha-\beta}f(t) + A\phi(t),
 \end{aligned}$$

for all $t \in \mathbb{R}$. This means that, ϕ is a (strong) solution to Eq (3.1). We recall that a function $u \in C(\mathbb{R}, X)$ is called a strong solution of (3.1) on \mathbb{R} if $u \in C(\mathbb{R}, D(A))$, the fractional derivative of u , $\partial^\alpha u$, exists and (3.1) holds for all $t \in \mathbb{R}$. If merely $u(t)$ belongs to X instead of the $D(A)$, then u is a mild solution to the Eq (3.1) according to Definition 8. As consequence of the above computation we have the following result.

Theorem 9. *Let $1 \leq \beta < \alpha < 2$ and $\omega < 0$. Assume that A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)}{2}\pi$. Then for each $f \in \mathcal{N}(X)$ there is a unique mild solution $u \in \mathcal{N}(X)$ of Eq (3.1) which is given by*

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds, \quad t \in \mathbb{R}.$$

Proof. By Theorem 5, the operator A generates a resolvent family $\{S_{\alpha,\beta}(t)\}_{t \geq 0}$ and by [28, Corollary 3.9] the function $t \mapsto S_{\alpha,\beta}(t)$ is uniformly 1-integrable. By Theorem 7 the function $u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds$ belongs to $\mathcal{N}(X)$ and it is the mild solution to (3.1). \square

Next, we consider the semilinear Eq (1.1).

Definition 10. *A function $u \in C(\mathbb{R}, X)$ is called a mild solution to Eq (1.1) if the function $s \mapsto S_{\alpha,\beta}(t-s)f(s, u(s))$ is integrable on $(-\infty, t)$ for each $t \in \mathbb{R}$ and*

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in \mathbb{R}. \quad (3.6)$$

Theorem 11. Let $1 \leq \beta < \alpha < 2$, $\omega < 0$ and A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)\pi}{2}$. If $f \in \mathcal{N}(\mathbb{R} \times X, X)$ satisfies

$$\|f(t, u) - f(t, v)\| \leq L\|u - v\|, \text{ for all } t \in \mathbb{R}, \text{ and } u, v \in X, \quad (3.7)$$

where $L < \frac{\alpha}{C}|\omega|^{\beta/\alpha} B\left(\frac{\beta}{\alpha}, 1 - \frac{\beta}{\alpha}\right)^{-1}$, and C is the constant given in Theorem 6, and $B(\cdot, \cdot)$ denotes the Beta function, then the Eq (1.1) has a unique mild solution $u \in \mathcal{N}(X)$.

Proof. Define the operator $F : \mathcal{N}(X) \rightarrow \mathcal{N}(X)$ by

$$(F\phi)(t) := \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s, \phi(s)) ds, \quad t \in \mathbb{R}. \quad (3.8)$$

By [28, Corollary 3.9] we have

$$\int_0^{\infty} \|S_{\alpha,\beta}(t)\| dt \leq \frac{C}{\alpha} |\omega|^{-\beta/\alpha} B\left(\frac{\beta}{\alpha}, 1 - \frac{\beta}{\alpha}\right) < \infty, \quad (3.9)$$

and [23, Theorems 3.3 and 4.1], F is well defined, that is, $F\phi \in \mathcal{N}(X)$ for all $\phi \in \mathcal{N}(X)$. For $\phi_1, \phi_2 \in \mathcal{N}(X)$ and $t \in \mathbb{R}$, by (3.9), we have:

$$\begin{aligned} \|(F\phi_1)(t) - (F\phi_2)(t)\| &\leq \int_{-\infty}^t \|S_{\alpha,\beta}(t-s)[f(s, \phi_1(s)) - f(s, \phi_2(s))]\| ds \\ &\leq \int_{-\infty}^t L \|S_{\alpha,\beta}(t-s)\| \cdot \|\phi_1(s) - \phi_2(s)\| ds \\ &\leq L \|\phi_1 - \phi_2\|_{\infty} \int_0^{\infty} \|S_{\alpha,\beta}(r)\| dr \\ &\leq \frac{LC}{\alpha} |\omega|^{-\beta/\alpha} B\left(\frac{\beta}{\alpha}, 1 - \frac{\beta}{\alpha}\right) \|\phi_1 - \phi_2\|_{\infty}. \end{aligned}$$

This proves that F is a contraction, so by the Banach fixed point theorem there exists a unique $u \in \mathcal{N}(X)$ such that $Fu = u$. \square

Theorem 12. Let $1 \leq \beta < \alpha < 2$, $\omega < 0$ and A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)\pi}{2}$. If $f \in \mathcal{N}(\mathbb{R} \times X, X)$ satisfies

$$\|f(t, u) - f(t, v)\| \leq \mathfrak{Q}(t)\|u - v\|, \text{ for all } t \in \mathbb{R}, \text{ and } u, v \in X,$$

where $\mathfrak{Q}(\cdot) \in L^1(\mathbb{R}, \mathbb{R}_+)$, then the Eq (1.1) admits a unique mild solution $u \in \mathcal{N}(X)$.

Proof. It easily follows by Theorem 6 that $\|S_{\alpha,\beta}(t)\| \leq \tilde{C} := \max\left\{C, \frac{C}{|\omega|}\right\}$. Define the operator F as (3.8). For $u, v \in \mathcal{N}(X)$ and $t \in \mathbb{R}$, we have

$$\begin{aligned} \|(Fu)(t) - (Fv)(t)\| &\leq \int_{-\infty}^t \|S_{\alpha,\beta}(t-s)[f(s, u(s)) - f(s, v(s))]\| ds \\ &\leq \tilde{C} \|u - v\|_{\infty} \int_0^{\infty} \mathfrak{Q}(t - \xi) d\xi \end{aligned}$$

$$= \tilde{C}\|u - v\|_{\infty} \int_{-\infty}^t \mathfrak{Q}(s) ds.$$

Generally, we have

$$\begin{aligned} \|(F^n u)(t) - (F^n v)(t)\| &\leq \|u - v\|_{\infty} \frac{(\tilde{C})^n}{(n-1)!} \left(\int_{-\infty}^t \mathfrak{Q}(s) \left(\int_{-\infty}^s \mathfrak{Q}(\xi) d\xi \right)^{n-1} ds \right) \\ &\leq \|u - v\|_{\infty} \frac{(\tilde{C})^n}{n!} \left(\int_{-\infty}^t \mathfrak{Q}(s) ds \right)^n \\ &\leq \|u - v\|_{\infty} \frac{(\|\mathfrak{Q}\|_1 \tilde{C})^n}{n!}. \end{aligned}$$

Since $\frac{(\|\mathfrak{Q}\|_1 \tilde{C})^n}{n!} < 1$ for sufficiently large n , by the contraction principle F admits a unique fixed point $u \in \mathcal{N}(X)$. \square

4. Mild solutions to Eq (1.2) with nonlocal conditions

Assume that A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)\pi}{2}$. By Theorem 5 the operator A generates a resolvent family $\{S_{\alpha,\beta}(t)\}_{t \geq 0}$. If $h : C(I, X) \rightarrow X$ is a continuous function, $f(0, u(0)) = 0$ and $u_1 \in X$, then it is well known that the mild solution to problem

$$\begin{cases} \partial_t^\alpha u(t) &= Au(t) + \partial_t^{\alpha-\beta} f(t, u(t)), & 0 \leq t \leq T \\ u(0) &= 0, \\ u'(0) + h(u) &= u_1, \end{cases} \quad (4.1)$$

is given by means of the variation-of-constant formula

$$u(t) = S_{\alpha,2}(t)[u_1 - h(u)] + \int_0^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in [0, T].$$

We assume the following

- **H1.** The function f satisfies the Carathéodory condition, that is $f(\cdot, u)$ is strongly measurable for each $u \in X$ and $f(t, \cdot)$ is continuous for each $t \in I := [0, T]$.
- **H2.** There exists a continuous function $\mu : I \rightarrow \mathbb{R}_+$ such that

$$\|f(t, u)\| \leq \mu(t)\|u\|, \quad \forall t \in I, u \in C(I, X)$$

and $f(0, u(0)) = 0$.

- **H3.** The function $h : C(I, X) \rightarrow X$ is continuous and there exists $L_h > 0$ such that

$$\|h(u) - h(v)\| < L_h\|u - v\|, \quad \forall u, v \in C(I, X).$$

- **H4.** The set $\mathcal{K} = \{S_{\alpha,\beta}(t-s)f(s, u(s)) : u \in C(I, X), 0 \leq s \leq t\}$ is relatively compact for each $t \in I$.

Proposition 13. Let $1 < \alpha < 2$ and $1 < \beta \leq 2$ such that $\alpha - \beta + 1 > 0$. If A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)}{2}\pi$, where $\omega < 0$, then the function $t \mapsto S_{\alpha,\beta}(t)$ is continuous in $\mathcal{B}(X)$ for all $t > 0$.

Proof. It proof follows similarly to [30, Proposition 11]. We omit the details. \square

We recall the following results.

Lemma 14 (Mazur's Theorem). If K is a compact subset of a Banach space X , then its convex closure $\text{conv}(K)$ is compact.

Lemma 15 (Leray-Schauder Alternative Theorem). Let C be a convex subset of a Banach space X . Suppose that $0 \in C$. If $F : C \rightarrow C$ is a completely continuous map, then either F has a fixed point, or the set $\{x \in C : x = \lambda F(x), 0 < \lambda < 1\}$ is unbounded.

Lemma 16 (Krasnoselskii Theorem). Let C be a closed convex and nonempty subset of a Banach space X . Let Q_1 and Q_2 be two operators such that

- i) If $u, v \in C$, then $Q_1u + Q_2v \in C$.
- ii) Q_1 is a mapping contraction.
- iii) Q_2 is compact and continuous.

Then, there exists $z \in C$ such that $z = Q_1z + Q_2z$.

We have the following existence theorem.

Theorem 17. Let $1 < \alpha < 2$ and $1 < \beta < 2$ such that $\alpha - \beta + 1 > 0$. Assume that A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)}{2}\pi$, where $\omega < 0$. Under assumptions H1-H4, the problem (4.1) has at least one mild solution.

Proof. By Theorem 5, the operator A generates a resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$. By the uniqueness of the Laplace transform we have $S_{\alpha,2}(t) = (g_1 * S_{\alpha,1})(t)$ and $S_{\alpha,\beta}(t) = (g_{\beta-1} * S_{\alpha,1})(t)$ for all $t \geq 0$. Moreover, by Theorem 6 there exists a constant $M > 0$ such that $\|S_{\alpha,2}(t)\| \leq M$ and $\|S_{\alpha,\beta}(t)\| \leq M$ for all $t \geq 0$. Now, we define the operator $\Gamma : C(I, X) \rightarrow C(I, X)$ by

$$(\Gamma u)(t) := S_{\alpha,2}(t)[u_1 - h(u)] + \int_0^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in [0, T].$$

Let $B_r := \{u \in C(I, X) : \|u\| \leq r\}$, where $r > 0$. We shall prove that Γ has at least one fixed point by the Leray-Schauder fixed point theorem. We will consider several steps in the proof.

Step 1. The operator Γ sends bounded sets of $C(I, X)$ into bounded sets of $C(I, X)$. In fact, take $u \in B_r$ and $G := \sup_{u \in B_r} \|h(u)\|$. Then

$$\begin{aligned} \|\Gamma u(t)\| &\leq \|S_{\alpha,2}(t)\|(\|u_1\| + \|h(u)\|) + \int_0^t \|S_{\alpha,\beta}(t-s)\| \|f(s, u(s))\| ds \\ &\leq M(\|u_1\| + G) + M \int_0^t \mu(s) \|u(s)\| ds \\ &\leq M(\|u_1\| + G) + Mr \int_0^t \mu(s) ds \\ &\leq M(\|u_1\| + G) + Mr \|\mu\|_{\infty} T := R. \end{aligned}$$

Therefore $\Gamma B_r \subset B_R$.

Step 2. Γ is a continuous operator.

Let $u_n, u \in B_r$ such that $u_n \rightarrow u$ in $C(I, X)$. Then we have

$$\begin{aligned} \|\Gamma u_n(t) - \Gamma u(t)\| &\leq \|S_{\alpha,2}(t)\|(\|h(u_n) - h(u)\|) + \int_0^t \|S_{\alpha,\beta}(t-s)\| \|f(s, u_n(s)) - f(s, u(s))\| ds \\ &\leq ML_h \|u_n - u\| + M \int_0^t \|f(s, u_n(s)) - f(s, u(s))\| ds \\ &\leq ML_h \|u_n - u\| + M \int_0^t \mu(s)(\|u_n(s)\| + \|u(s)\|) ds \\ &\leq ML_h \|u_n - u\| + 2rM \int_0^t \mu(s) ds. \end{aligned}$$

We notice that the function $s \mapsto \mu(s)$ is integrable on I . By the Lebesgue's Dominated Convergence Theorem, $\int_0^t \|f(s, u_n(s)) - f(s, u(s))\| ds \rightarrow 0$ as $n \rightarrow \infty$. Since $u_n \rightarrow u$ we obtain that Γ is continuous in $C(I, X)$.

Step 3 The operator Γ sends bounded sets of $C(I, X)$ into equicontinuous sets of $C(I, X)$.

In fact, let $u \in B_r$, with $r > 0$ and take $t_1, t_2 \in I$ with $t_2 < t_1$. Then we have

$$\begin{aligned} \|\Gamma u(t_1) - \Gamma u(t_2)\| &\leq \|(S_{\alpha,2}(t_1) - S_{\alpha,2}(t_2))(u_1 - h(u))\| + \int_{t_2}^{t_1} \|S_{\alpha,\beta}(t_1 - s)f(s, u(s))\| ds \\ &\quad + \int_0^{t_2} \|(S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s))f(s, u(s))\| ds \\ &:= I_1 + I_2 + I_3. \end{aligned}$$

Observe that

$$I_1 \leq \|(S_{\alpha,2}(t_1) - S_{\alpha,2}(t_2))\| \|u_1 - h(u)\|.$$

Using the norm continuity of $t \mapsto S_{\alpha,2}(t)$ (see Proposition 13) we obtain that $\lim_{t_1 \rightarrow t_2} I_1 = 0$.

On the other hand,

$$I_2 \leq M \int_{t_2}^{t_1} \mu(s) \|u(s)\| ds \leq rM \|\mu\|_\infty (t_1 - t_2),$$

and therefore $\lim_{t_1 \rightarrow t_2} I_2 = 0$. Finally, for I_3 we have

$$\begin{aligned} I_3 &\leq \int_0^{t_2} \|S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s)\| \|f(s, u(s))\| ds \\ &\leq \int_0^{t_2} \|S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s)\| \mu(s) \|u(s)\| ds \\ &\leq r \int_0^{t_2} \|S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s)\| \mu(s) ds. \end{aligned}$$

Since

$$\|S_{\alpha,\beta}(t_1 - \cdot) - S_{\alpha,\beta}(t_2 - \cdot)\| \mu(\cdot) \leq 2M \mu(\cdot) \in L^1(I, \mathbb{R}),$$

and $S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s) \rightarrow 0$ in $\mathcal{B}(X)$, as $t_1 \rightarrow t_2$ (see Proposition 13) we obtain by the Lebesgue's dominated convergence theorem that $\lim_{t_1 \rightarrow t_2} I_3 = 0$. The proof of the claim is finished.

Step 4. The function Γ maps B_r into relatively compact sets in X .

The hypothesis and Lemma 14 imply that $\text{conv}(\mathcal{K})$ is compact. Moreover, for $u \in B_r$, by the Mean-Value Theorem for the Bochner integral (see [15, Corollary 8, p. 48]), we get

$$\Gamma(u(t)) \in \overline{t \text{conv}(\mathcal{K})},$$

for all $t \in [0, T]$. Thus the set $\overline{\{\Gamma u(t); u \in B_r\}}$ is compact in X for every $t \in [0, T]$.

We conclude from Steps 1–4, that Γ is continuous and compact by the Arzela-Ascoli's theorem, which means that the function Γ is completely continuous.

Step 5. The set $\Omega := \{u \in B_r : u = \lambda \Gamma u, 0 < \lambda < 1\}$ is bounded. In fact, since $0 \in \Omega$ we obtain that $\Omega \neq \emptyset$. For $u \in \Omega$ we have

$$\begin{aligned} \|u(t)\| &\leq \lambda[M(\|u_1\| + \|h(u)\|) + M \int_0^t \|f(s, u(s))\| ds] \\ &\leq \lambda[M(\|u_1\| + G) + M \int_0^t \mu(s)\|u(s)\| ds] \\ &\leq [M(\|u_1\| + G) + Mr\|\mu\|_\infty T], \end{aligned}$$

for all $t \in [0, T]$, which means that Ω is a bounded set.

Therefore, by Lemma 15 we conclude that Γ has a fixed point, and the proof of the Theorem is finished. \square

The same method of proof can be used to prove the next result. We omit the details.

Theorem 18. *Let $1 < \alpha < 2$. Assume that A generates the resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$. Under assumptions H1–H4, the problem (4.1) has at least one mild solution.*

Now, we consider the problem

$$\begin{cases} \partial_t^\alpha u(t) &= Au(t) + \partial_t^{\alpha-\beta} f(t, u(t)), & 0 \leq t \leq T \\ u(0) + g(u) &= u_0, \\ u'(0) + h(u) &= u_1, \end{cases} \quad (4.2)$$

where $g, h : C(I, X) \rightarrow X$ are continuous, $f(0, u(0)) = 0$ and $u_0, u_1 \in X$. By (2.2) in Theorem 6, there exists a constant $M > 0$ such that

$$\|S_{\alpha,1}(t)\| \leq \frac{M}{1 + |\omega|t^\alpha}, \quad \|S_{\alpha,2}(t)\| \leq \frac{Mt}{1 + |\omega|t^\alpha}, \quad \|S_{\alpha,\beta}(t)\| \leq \frac{Mt^{\beta-1}}{1 + |\omega|t^\alpha}, \quad t \geq 0. \quad (4.3)$$

Thus

$$\|S_{\alpha,1}(t)\| \leq M, \quad \|S_{\alpha,2}(t)\| \leq MT, \quad \|S_{\alpha,\beta}(t)\| \leq MT^{\beta-1}, \quad t \in [0, T]. \quad (4.4)$$

Under the same assumptions H1–H3 and

- **H3'**. The function $g : C(I, X) \rightarrow X$ is continuous and there exists $L_g > 0$ such that

$$\|g(u) - g(v)\| < L_g \|u - v\|, \quad \forall u, v \in C(I, X).$$

we have the following result.

Theorem 19. Let $1 < \alpha < 2$ and $1 < \beta < 2$ such that $\alpha - \beta + 1 > 0$. Assume that A is an ω -sectorial operator of angle $\theta = \frac{(\alpha-1)}{2}\pi$, where $\omega < 0$. Suppose that $M\|\mu\|_{\infty}T^{\beta} < 1$ and $M(L_g + TL_h) < 1$, where M is the constant in (4.4). Assume that $(\lambda^{\alpha} - A)^{-1}$ is compact for all $\lambda > \nu^{1/\alpha}$, where ν is a positive constant. Under assumptions H1–H3 and H3', the problem (4.2) has at least one mild solution.

Proof. By Theorem 5, the operator A generates the resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$, and $S_{\alpha,2}(t) = (g_1 * S_{\alpha,1})(t)$ and $S_{\alpha,\beta}(t) = (g_{\beta-1} * S_{\alpha,1})(t)$ for all $t \geq 0$. Then, the mild solution to problem (4.2) is given by

$$u(t) = S_{\alpha,1}(t)[u_0 - g(u)] + S_{\alpha,2}(t)[u_1 - h(u)] + \int_0^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in [0, T].$$

Let $B_r := \{u \in C(I, X) : \|u\| \leq r\}$, where

$$r := \frac{M(\|u_0\| + \|g(u)\|) + MT(\|u_1\| + \|h(u)\|)}{1 - M\|\mu\|_{\infty}T^{\beta}}.$$

On B_r we define the operators Γ_1, Γ_2 by

$$\begin{aligned} (\Gamma_1 u)(t) &:= S_{\alpha,1}(t)[u_0 - g(u)] + S_{\alpha,2}(t)(u_1 - h(u)) \quad t \in [0, T] \\ (\Gamma_2 u)(t) &:= \int_0^t S_{\alpha,\beta}(t-s)f(s, u(s))ds, \quad t \in [0, T], \end{aligned}$$

where $u \in B_r$. We claim that $\Gamma := \Gamma_1 + \Gamma_2$ has at least one fixed. To prove this, we will consider several steps.

Step 1. We claim that if $u, v \in B_r$, then $\Gamma_1 u + \Gamma_2 v \in B_r$. In fact,

$$\begin{aligned} \|(\Gamma_1 u)(t) + (\Gamma_2 v)(t)\| &\leq \\ &\leq \|S_{\alpha,1}(t)\| \|u_0 - g(u)\| + \|S_{\alpha,2}(t)\| \|u_1 - h(u)\| + \int_0^t \|S_{\alpha,\beta}(t-s)\| \|f(s, v(s))\| ds \\ &\leq M(\|u_0\| + \|g(u)\|) + MT(\|u_1\| + \|h(u)\|) + M \int_0^t (t-s)^{\beta-1} \mu(s) \|v(s)\| ds \\ &\leq M(\|u_0\| + \|g(u)\|) + MT(\|u_1\| + \|h(u)\|) + MT^{\beta} \|\mu\|_{\infty} r = r. \end{aligned}$$

Thus $\Gamma_1 u + \Gamma_2 v \in B_r$ for all $u, v \in B_r$.

Step 2. Γ_1 is a contraction on B_r . In fact, if $u, v \in B_r$, then

$$\|\Gamma_1 u(t) - \Gamma_1 v(t)\| \leq \|S_{\alpha,1}(t)\| \|g(u) - g(v)\| + \|S_{\alpha,2}(t)\| \|h(u) - h(v)\| \leq (ML_g + MTL_h) \|u - v\|$$

Since $M(L_g + TL_h) < 1$, we get that Γ_1 is a contraction.

Step 3. Γ_2 is completely continuous.

Firstly, we prove that Γ_2 is a continuous operator on B_r . Let $u_n, u \in B_r$ such that $u_n \rightarrow u$ in B_r . We notice that by (4.3)

$$\|\Gamma_2 u_n(t) - \Gamma_2 u(t)\| \leq \int_0^t \|S_{\alpha,\beta}(t-s)\| \|f(s, u_n(s)) - f(s, u(s))\| ds \leq 2MrT^{\beta} \int_0^t \mu(s) ds.$$

Moreover, the function $s \mapsto \mu(s)$ is integrable on $[0, T]$. The Lebesgue's Dominated Convergence Theorem implies that $\int_0^t \|f(s, u_n(s)) - f(s, u(s))\| ds \rightarrow 0$ as $n \rightarrow \infty$. Since $u_n \rightarrow u$ we obtain that Γ_2 is continuous in B_r .

Now, we prove that $\{\Gamma_2 u : u \in B_r\}$ is a relatively compact set. In fact, by the Ascoli-Arzelà theorem we only need to prove that the family $\{\Gamma_2 u : u \in B_r\}$ is uniformly bounded and equicontinuous, and the set $\{\Gamma_2 u(t) : u \in B_r\}$ is relatively compact in X for each $t \in [0, T]$. For each $u \in B_r$ we have $\|\Gamma_2 u\| \leq MT^\beta r \|\mu\|_\infty$, which implies that $\{\Gamma_2 u : u \in B_r\}$ is uniformly bounded.

Next, we prove the equicontinuity. For $u \in B_r$ and $0 \leq t_2 < t_1 \leq T$ we have

$$\begin{aligned} \|\Gamma_2 u(t_1) - \Gamma_2 u(t_2)\| &\leq \int_{t_2}^{t_1} \|S_{\alpha,\beta}(t_1 - s)f(s, u(s))\| ds \\ &+ \int_0^{t_2} \|(S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s))f(s, u(s))\| ds =: I_1 + I_2. \end{aligned}$$

Observe that for I_1 , by (4.3) we have $I_1 \leq MT^\beta \int_{t_2}^{t_1} \mu(s) \|u(s)\| ds \leq MT^\beta r \|\mu\|_\infty (t_1 - t_2)$, and thus $\lim_{t_1 \rightarrow t_2} I_1 = 0$. On the other hand, for I_2 we have

$$I_2 \leq \int_0^{t_2} \|S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s)\| \|f(s, u(s))\| ds \leq r \int_0^{t_2} \mu(s) \|S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s)\| ds.$$

By (4.4) we have $\mu(\cdot) \|S_{\alpha,\beta}(t_1 - \cdot) - S_{\alpha,\beta}(t_2 - \cdot)\| \leq 2T^{\beta-1} M \mu(\cdot) \in L^1([0, T], \mathbb{R})$, and by Proposition 13 the function $t \mapsto S_{\alpha,\beta}(t)$ is norm continuous. This implies that if $t_1 \rightarrow t_2$, then $S_{\alpha,\beta}(t_1 - s) - S_{\alpha,\beta}(t_2 - s) \rightarrow 0$ in $\mathcal{B}(X)$. By the Lebesgue's dominated convergence theorem we conclude that $\lim_{t_1 \rightarrow t_2} I_2 = 0$. Therefore, $\{\Gamma_2 u : u \in B_r\}$ is an equicontinuous family.

Finally, we prove that $H(t) := \{\Gamma_2 u(t) : u \in B_r\}$ is relatively compact in X for each $t \in [0, T]$. Clearly, $H(0)$ is relatively compact in X . Now, we take $t > 0$. For $0 < \varepsilon < t$ we define on B_r the operator

$$(\Gamma_2^\varepsilon u)(t) := \int_0^{t-\varepsilon} S_{\alpha,\beta}(t-s)f(s, u(s)) ds.$$

By [30, Theorem 14] we have that $S_{\alpha,\beta}(t)$ is a compact operator for all $t > 0$. Thus $\mathcal{K}_\varepsilon := \{S_{\alpha,\beta}(t-s)f(s, u(s)) : u \in B_r, 0 \leq s \leq t-\varepsilon\}$ is a compact set for all $\varepsilon > 0$. By Lemma 14, $\overline{\text{conv}(\mathcal{K}_\varepsilon)}$ is also a compact set. The Mean-Value Theorem for the Bochner integrals (see [15, Corollary 8, p. 48]), implies that $(\Gamma_2^\varepsilon u)(t) \in t \text{conv}(\mathcal{K}_\varepsilon)$, for all $t \in [0, T]$. Therefore, the set $H_\varepsilon(t) := \{(\Gamma_2^\varepsilon u)(t) : u \in B_r\}$ is relatively compact in X for all $\varepsilon > 0$. Since

$$\|(\Gamma_2 u)(t) - (\Gamma_2^\varepsilon u)(t)\| \leq \int_{t-\varepsilon}^t \|S_{\alpha,\beta}(t-s)f(s, u(s))\| ds \leq MT^{\beta-1} r \int_{t-\varepsilon}^t \mu(s) ds$$

and the function $s \mapsto \mu(s)$ belongs to $L^1([t-\varepsilon, t], \mathbb{R}_+)$ we conclude by the Lebesgue dominated convergence Theorem that $\lim_{\varepsilon \rightarrow 0} \|(\Gamma_2 u)(t) - (\Gamma_2^\varepsilon u)(t)\| = 0$. Therefore the set $\{\Gamma_2 u(t) : u \in B_r\}$ is relatively compact in X for each $t \in (0, T]$. The Ascoli-Arzelà theorem implies that the set $\{\Gamma_2 u : u \in B_r\}$ is relatively compact. We conclude that Γ_2 is a completely continuous operator. By Lemma 16 we have that $\Gamma = \Gamma_1 + \Gamma_2$ has a fixed point on B_r , and therefore the problem (4.2) has a mild solution. \square

5. Examples

Example 20.

On the Banach space $X = \mathbb{C}$, let A be the scalar operator $A = \varrho I$, where $\varrho \in \mathbb{R}$. Consider the multi-term fractional differential equation

$$\partial^\alpha u(t) = Au(t) + \partial^{\alpha-\beta} f(t), \quad t \in \mathbb{R}, \quad (5.1)$$

where $1 \leq \beta < \alpha < 2$ and $f(t)$ is the almost periodic function $f(t) = \sin(t) + \sin(\sqrt{2}t)$, see [14, p. 80]. By Theorem 9 the solution u to (5.1) is an almost periodic function, and it is given by

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds, \quad t \in \mathbb{R}.$$

where $S_{\alpha,\beta}(t) = t^{\beta-1}E_{\alpha,\beta}(\varrho t^\alpha)$. By Theorem 6, we can write

$$u(t) = \int_{-\infty}^t S_{\alpha,\beta}(t-s)f(s)ds = \sum_{k=0}^{\infty} \varrho^k \int_{-\infty}^t \frac{(t-s)^{\alpha k + \beta - 1}}{\Gamma(\alpha k + \beta)} f(s)ds.$$

Now, we notice that if $g(t) = e^{\mu t}$, where $\mu \in \mathbb{C}$ and $\delta > 0$, then

$$\frac{1}{\Gamma(\delta)} \int_{-\infty}^t (t-s)^{\delta-1} g(s)ds = \frac{\mu^{1-\delta}}{\Gamma(\delta)} \int_{-\infty}^t [\mu(t-s)^{\delta-1}] e^{\mu s} ds = \frac{\mu^{-\delta}}{\Gamma(\delta)} e^{\mu t} \int_0^{\infty} r^{\delta-1} e^{-r} dr = \mu^{-\delta} e^{\mu t},$$

and therefore, for $h(t) = \sin(at) = \frac{e^{ait} - e^{-ait}}{2i}$, where $a > 0$, we have

$$\frac{1}{\Gamma(\delta)} \int_{-\infty}^t (t-s)^{\delta-1} h(s)ds = a^{-\delta} \sin\left(at - \frac{\pi}{2}\delta\right).$$

This implies that

$$u(t) = \sum_{k=0}^{\infty} \varrho^k \left[\sin\left(t - \frac{\pi}{2}(\alpha k + \beta)\right) + \frac{1}{\sqrt{2}^{\alpha k + \beta}} \sin\left(\sqrt{2}t - \frac{\pi}{2}(\alpha k + \beta)\right) \right].$$

In Figure 1, we have the solution u for (5.1) for $\varrho = -1$ and $\alpha = 1.5, \beta = 1.3$ on the interval $[-30, 30]$.

Example 21.

Consider the following partial differential equation with fractional temporal derivatives

$$\left\{ \begin{array}{l} \partial_t^\alpha u(t, x) = \frac{\partial^2}{\partial x^2} u(t, x) + \partial_t^{\alpha-\beta} \sin(u(t, x)), \quad (t, x) \in [0, T] \times \mathbb{R} \\ u(0, x) = 0, \quad x \in \mathbb{R} \\ u'(0, x) + \sum_{i=1}^n c_i u(t_i, x) = u_1(x), \quad x \in \mathbb{R} \end{array} \right. \quad (5.2)$$

where $1 < \alpha, \beta < 2$, $0 \leq t_1 < \dots < t_n \leq T$, $u_1 \in L^2(\mathbb{R})$, and c_i are real constants.




Figure 1. Solution $u(t)$ for (5.1) on the interval $[-30, 30]$.

On the Banach space $X = L^2(\mathbb{R})$, let A be the second order operator $Av = v''$ with domain $D(A) = W^{2,2}(\mathbb{R})$. By [36, Example 1.2.2, p. 3063], A generates a cosine family $\{S_{2,1}(t)\}_{t \in \mathbb{R}}$ on X , and by the Subordination Principle [32, Corollary 3.3], A is the generator of the resolvent family $\{S_{\alpha,1}(t)\}_{t \geq 0}$ given by

$$S_{\alpha,1}(t)x := \int_0^\infty \psi_{\frac{\alpha}{2}, 1-\frac{\alpha}{2}}(t, s) S_{2,1}(s)x ds, \quad t \geq 0, x \in X, \quad (5.3)$$

where $\psi_{\frac{\alpha}{2}, 1-\frac{\alpha}{2}}$ is the Wright type function defined by

$$\begin{aligned} \psi_{\frac{\alpha}{2}, 1-\frac{\alpha}{2}}(t, s) = & \frac{1}{\pi} \int_0^\infty \rho^{\frac{\alpha}{2}-1} e^{-s\rho^{\frac{\alpha}{2}} \cos \frac{\alpha}{2}(\pi-\theta) - t\rho \cos \theta} \\ & \times \sin \left(t\rho \sin \theta - s\rho^{\frac{\alpha}{2}} \sin \frac{\alpha}{2}(\pi-\theta) + \frac{\alpha}{2}(\pi-\theta) \right) d\rho, \end{aligned}$$

for $\theta \in (\pi - \frac{2}{\alpha}, \pi/2)$. Define,

$$\begin{aligned} u(t)x &= u(t, x) \\ f(t, u(t))(x) &= \sin(u(t, x)) \\ h(u)(x) &= \sum_{i=1}^n c_i u(t_i, x). \end{aligned}$$

Then, (5.2) can be reformulated as the abstract problem (4.1). Moreover, an easy computation shows that the hypotheses **H1–H3** hold with $\mu(t) = 1$ and $L_h = \sum_{i=1}^n |c_i|$. Since

$$S_{\alpha,\beta}(t) = (g_{\beta-1} * S_{\alpha,1})(t),$$

we obtain by (2.2) that the set $\mathcal{K} = \{S_{\alpha,\beta}(t-s) \sin(s, u(s)) : u \in C(I, X), 0 \leq s \leq t\}$ is relatively compact for each $t \in I$, and therefore **H4** holds. We conclude, by Theorem 18, that the problem (4.1) has at least one mild solution u .

6. Conclusions

In this paper, some sufficient conditions are established for the existence of mild and bounded solutions to a multi-term fractional differential equation. Under the theory of fractional resolvent families in Banach spaces, the mild solutions are given in term of these operator families. The main results are given for the Caputo and Weyl fractional derivatives.

Acknowledgements

The authors thank the anonymous referees for reading the manuscript and for making suggestions that have improved the previous version of this paper. This work was partially supported by NSF of Shaanxi Province (2020JM-183).

Conflict of interest

The authors declare no conflict of interest.

References

1. M. Ali, S. A. Malik, An inverse problem for a family of two parameters time fractional diffusion equations with nonlocal boundary conditions, *Math. Methods Appl. Sci.*, **40** (2017), 7737–7748.
2. E. Alvarez-Pardo, C. Lizama, Mild solutions for multi-term time-fractional differential equations with nonlocal initial conditions, *Electron. J. Differ. Equations*, **39** (2014), 1–10.
3. B. de Andrade, C. Cuevas, H. Soto, On fractional heat equations with non-local initial conditions, *Proc. Edinburgh Math. Soc.*, **59** (2016), 65–76.
4. D. Araya, C. Lizama, Almost automorphic mild solutions to fractional differential equations, *Nonlinear Anal.: Theory Methods Appl.*, **69** (2008), 3692–3705.
5. W. Arendt, C. Batty, M. Hieber, F. Neubrander, *Vector-Valued Laplace Transforms and Cauchy Problems*, Birkhäuser Verlag, Basel, 2011.
6. L. Byszewski, Theorems about the existence and uniqueness of solutions of a semilinear evolution nonlocal Cauchy problem, *J. Math. Anal. Appl.*, **162** (1991), 494–505.
7. A. Caicedo, C. Cuevas, G. M. Mophou, G. N'Guérékata, Asymptotic behavior of solutions of some semilinear functional differential and integro-differential equations with infinite delay in Banach spaces, *J. Franklin Inst.*, **349** (2012), 1–24.
8. J. Cao, Q. Yang, Z. Huang, Existence of anti-periodic mild solutions for a class of semilinear fractional differential equations, *Commun. Nonlinear Sci. Numer. Simul.*, **17** (2012), 277–283.
9. P. M. de Carvalho-Neto, G. Planas, Mild solutions to the time fractional Navier-Stokes equations in \mathbb{R}^N , *J. Differ. Equations*, **259** (2015), 2948–2980.
10. P. Chen, X. Zhang, Y. Li, Existence and approximate controllability of fractional evolution equations with nonlocal conditions via resolvent operators, *Fract. Calculus Appl. Anal.*, **23** (2020), 268–291.

11. E. Cuesta, Asymptotic behaviour of the solutions of fractional integro-differential equations and some time discretizations, *Conference Publications, American Institute of Mathematical Sciences*, **2007** (2007), 277–285.
12. C. Cuevas, C. Lizama, Almost automorphic solutions to a class of semilinear fractional differential equations, *Appl. Math. Lett.*, **21** (2008), 1315–1319.
13. C. Cuevas, J. C. de Souza, S -asymptotically ω -periodic solutions of semilinear fractional integro-differential equations, *Appl. Math. Lett.*, **22** (2009), 865–870.
14. T. Diagana, *Almost Automorphic Type and Almost Periodic Type Functions in Abstract Spaces*, Springer, New York, 2013.
15. J. Diestel, J. J. Uhl, *Vector Measures*, American Mathematical Society, Providence, 1977.
16. S. D. Eidelman, A. N. Kochubei, Cauchy problem for fractional diffusion equations, *J. Differ. Equations*, **199** (2004), 211–255.
17. K. J. Engel, R. Nagel, One-parameter semigroups for linear evolution equations, *Semigroup forum, Springer-Verlag*, **63** (2001), 278–280.
18. M. Haase, *The Functional Calculus for Sectorial Operators*, Birkäuser Verlag, Basel, 2006.
19. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier Science B.V., Amsterdam, 2006.
20. K. Li, J. Peng, J. Jia, Cauchy problems for fractional differential equations with Riemann-Liouville fractional derivatives, *J. Funct. Anal.*, **263** (2012), 476–510.
21. C. Lizama, Regularized solutions for abstract Volterra equations, *J. Math. Anal. Appl.*, **243** (2000), 278–292.
22. C. Lizama, Solutions of two-term time fractional order differential equations with nonlocal initial conditions, *Electron. J. Qualitative Theory Differ. Equations*, **82** (2012), 1–9.
23. C. Lizama, G. M. N’Guérékata, Bounded mild solutions for semilinear integro-differential equations in Banach spaces, *Integr. Equations Oper. Theory*, **68** (2010), 207–227.
24. C. Lizama, G. M. N’Guérékata, Mild solutions for abstract fractional differential equations, *Appl. Anal.*, **92** (2013), 1731–1754.
25. K. S. Miller, B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, Wiley, New York, 1993.
26. G. M. Mophou, G. M. N’Guérékata, Existence of the mild solution for some fractional differential equations with nonlocal conditions, *Semigroup Forum, Springer-Verlag*, **79** (2009), 315–322.
27. G. M. N’Guérékata, *Topics in Almost Automorphy*, Springer Verlag, New York, 2005.
28. R. Ponce, Asymptotic behavior of mild solutions to fractional Cauchy problems in Banach spaces, *Appl. Math. Lett.*, **105** (2020), 106322.
29. R. Ponce, Bounded mild solutions to fractional integro-differential equations in Banach spaces, *Semigroup Forum, Springer US*, **87** (2013), 377–392.
30. R. Ponce, Existence of mild solutions to nonlocal fractional Cauchy problems via compactness, *Abstr. Appl. Anal.*, **2016** (2016), 4567092.

31. R. Ponce, Hölder continuous solutions for fractional differential equations and maximal regularity, *J. Differ. Equations*, **255** (2013), 3284–3304.
32. R. Ponce, Subordination principle for fractional diffusion-wave equations of Sobolev type, *Fract. Calculus Appl. Anal.*, **23** (2020), 427–449.
33. X. B. Shu, Q. Wang, The existence and uniqueness of mild solutions for fractional differential equations with nonlocal conditions of order $1 < \alpha < 2$, *Comput. Math. Appl.*, **64** (2012), 2100–2110.
34. L. Shu, X. B. Shu, J. Mao, Approximate controllability and existence of mild solutions for Riemann-Liouville fractional stochastic evolution equations with nonlocal conditions of order $1 < \alpha < 2$, *Fract. Calculus Appl. Anal.*, **22** (2019), 1086–1112.
35. B. T. Torebek, R. Tapdigoglu, Some inverse problems for the nonlocal heat equation with Caputo fractional derivative, *Math. Methods Appl. Sci.*, **40** (2017), 6468–6479.
36. V. V. Vasilev, S. I. Piskarev, Differential equations in Banach spaces. II. Theory of cosine operator functions, *J. Math. Sci.*, **122** (2004), 3055–3174.
37. G. Wang, X. Ren, Z. Bai, W. Hou, Radial symmetry of standing waves for nonlinear fractional Hardy-Schrödinger equation, *Appl. Math. Lett.*, **96** (2019), 131–137.
38. R. N. Wang, T. J. Xiao, J. Liang, A note on the fractional Cauchy problems with nonlocal initial conditions, *Appl. Math. Lett.*, **24** (2011), 1435–1442.
39. A. Wiman, Über den Fundamentalsatz in der Theorie der Funktionen $E^\alpha(x)$, *Acta Math.*, **29** (1905), 191–201.
40. A. Wiman, Über die Nullstellen der Funktionen $E^\alpha(x)$, *Acta Math.*, **29** (1905), 217–234.
41. X. J. Yang, *General Fractional Derivatives: Theory, Methods and Applications*, CRC Press, New York, 2019.
42. X. J. Yang, F. Gao, Y. Ju, *General Fractional Derivatives with Applications in Viscoelasticity*, Academic Press, 2020.
43. X. J. Yang, New insight into the Fourier-like and Darcy-like models in porous medium, *Thermal Sci.*, **24** (2020), 3847–385.
44. R. Zacher, A weak Harnack inequality for fractional differential equations, *J. Integral Equations Appl.*, **19** (2007), 209–232.
45. Y. Zhou, *Basic Theory of Fractional Differential Equations*, World Scientific, 2014.



AIMS Press

©2021 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)