



*Research article*

## A basic study of a fractional integral operator with extended Mittag-Leffler kernel

Gauhar Rahman<sup>1,\*</sup>, Iyad Suwan<sup>2</sup>, Kottakkaran Sooppy Nisar<sup>3</sup>, Thabet Abdeljawad<sup>4,5,\*</sup>, Muhammad Samraiz<sup>6</sup> and Asad Ali<sup>1</sup>

<sup>1</sup> Department of Mathematics and Statistics, Hazara University, Mansehra, Pakistan

<sup>2</sup> Department of Mathematics and Statistics, The Arab American University, P.O. Box 240, 13 Zababdeh, Jenin, Palestine

<sup>3</sup> Department of Mathematics, College of Arts and Sciences, Wadi Aldawser 11991, Prince Sattam bin Abdulaziz University, Saudi Arabia

<sup>4</sup> Department of Mathematics and General Sciences, Prince Sultan University, Riyadh, KSA

<sup>5</sup> Department of Medical Research, China Medical University, Taichung 40402, Taiwan

<sup>6</sup> Samraiz-Department of Mathematics, University of Sargodha, P.O. Box 40100, Sargodha, Pakistan

\* **Correspondence:** Email: [gauhar55uom@gmail.com](mailto:gauhar55uom@gmail.com), [tabdeljawad@psu.edu.sa](mailto:tabdeljawad@psu.edu.sa).

**Abstract:** In this present paper, the basic properties of an extended Mittag-Leffler function are studied. We present some fractional integral and differential formulas of an extended Mittag-Leffler function. In addition, we introduce a new extension of Prabhakar type fractional integrals with an extended Mittag-Leffler function in the kernel. Also, we present certain basic properties of the generalized Prabhakar type fractional integrals.

**Keywords:** Mittag-Leffler function; fractional integral; Prabhakar fractional integral

**Mathematics Subject Classification:** 26A33, 26D10, 26D53, 05A30

### 1. Introduction

In the last few years, many mathematical models were developed for various real-world problems by utilizing the field of fractional calculus with boundary conditions. Therefore, this field has received more attention in a variety of fields in diverse domains see [13, 14, 18, 32, 33]. In the development of fractional calculus, many researchers focused on developing certain new fractional integral operators and their applications in diverse fields. In practical applications, certain various types of fractional operators such as Riemann-Liouville, Caputo, Riesz [11, 35] and Hilfer [12] fractional operators are

introduced. Freshly, the researchers have studied certain new fractional integral and derivative operators and their possible applications in various disciplines of sciences. Very recently Khalil et al. [9] have introduced the notion of fractional conformable derivative (FCD) operators with some shortcomings. Abdeljawad [1] investigated the properties of the fractional conformable derivative operators. In [8], Jarad et al. introduced the fractional conformable integral and derivative operators. Anderson and Unles [4] developed the idea of conformable derivative by employing local proportional derivatives. Abdeljawad and Baleanu [2] investigated certain monotonicity results for fractional difference operators with discrete exponential kernels. Abdeljawad and Baleanu [3] have established fractional derivative operators with exponential kernel and their discrete versions. In [5], Atangana and Baleanu defined a new fractional derivative operator with the non-local and non-singular kernel. Caputo and Fabrizio [6] defined fractional derivative without a singular kernel. Certain properties of fractional derivative without a singular kernel can be found in the work of Losada and Nieto [10]. In [7], Jarad et al. defined generalized fractional derivatives generated by a class of local proportional derivatives.

On the other side, the researchers studied the fractional operators and have investigated and studied fractional integral inequalities to examine the existence and uniqueness of such type of fractional boundary value problems [46–52].

The special functions such as Mittag-Leffler function and its extensions appear as solution of fractional order differential and integral equations. Various interesting applications of such functions can be found in the study of the telegraph equations, kinetic equation, Levy flights, random walks, complex system, non-equilibrium statistical mechanics, super diffuse transport and quantum mechanics. The interested readers may refer to the work [15, 26–28] and the references cited therein.

## 2. Preliminaries

In this section, we present some basic and well known results.

In [53], the Gosta Mittag-Leffler function  $E_\rho(z)$  is defined by

$$E_\rho(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\rho n + 1)}, \quad (z \in \mathbb{C}; \Re(\rho) > 0). \quad (2.1)$$

The Wiman Mittag-Leffler function  $E_{\rho,\varphi}(z)$  is defined in [54] by

$$E_{\rho,\varphi}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\rho n + \varphi)}, \quad (z, \varphi \in \mathbb{C}; \Re(\rho) > 0). \quad (2.2)$$

The researchers extensively studied  $E_\rho(z)$  and  $E_{\rho,\varphi}(z)$  [17, 20–22, 24, 25, 34]. The numerical investigation of such functions in the complex plane can be found in the work of [23, 36]. Prabhakar [31] has proposed the following generalization of Mittag-Leffler function by

$$E_{\rho,\varphi}^\delta(z) = \sum_{n=0}^{\infty} \frac{(\delta)_n}{\Gamma(\rho n + \varphi)} \frac{z^n}{n!}, \quad (z, \varphi \in \mathbb{C}; \Re(\rho) > 0), \quad (2.3)$$

where  $(\delta)_n$  denotes the well-known Pochhammer symbol.

Rahman et al. [29] recently defined an extension of Mittag-Leffler function by

$$E_{\rho, \varphi; p, \nu}^{\delta}(z) = \sum_{n=0}^{\infty} \frac{(\delta; p, \nu)_n z^n}{\Gamma(\rho n + \varphi) n!}, \quad (z, \rho, \varphi \in \mathbb{C}; \Re(\rho) > 0, \Re(\varphi) > 0, \Re(p) > 0, p, \nu \geq 0), \quad (2.4)$$

where  $(\delta; p, \nu)_n$  is an extended Pochhammer symbol recently defined by Srivastava et al. [44] by

$$(\delta; p, \nu)_n = \begin{cases} \frac{\Gamma(\delta+n;p)}{\Gamma(\delta)}, & (\Re(p) > 0, p, \nu \geq 0, \delta \in \mathbb{C}), \\ (\delta; p)_n, & (\nu = 0, \delta \in \mathbb{C}). \end{cases} \quad (2.5)$$

An extension of gamma function  $\Gamma_{\nu}(\delta; p)$  is defined by

$$\Gamma_{\nu}(\delta; p) = \sqrt{\frac{2p}{\pi}} \int_0^{\infty} t^{\delta-\frac{3}{2}} e^{-t} K_{\nu+\frac{1}{2}}\left(\frac{p}{t}\right) dt,$$

where  $K_{\nu+\frac{1}{2}}\left(\frac{p}{t}\right)$  is the modified Bessel function of the first kind (see, e.g., Chaudhry and Zubair [16]). If we consider  $\nu = 0$  in (2.5), then we get the following extension of Pochhammer symbol defined by Srivastava et al. [45] as

$$(\delta; p)_n = \begin{cases} \frac{\Gamma(\delta+n;p)}{\Gamma(\delta)}, & (\Re(p) > 0, p \geq 0, \delta \in \mathbb{C}), \\ (\delta)_n, & (\delta \in \mathbb{C}), \end{cases} \quad (2.6)$$

where  $\Gamma(\delta; p)$  is defined by

$$\Gamma(\delta; p) = \int_0^{\infty} t^{\delta-1} e^{-t-\frac{p}{t}} dt.$$

Similarly, if we consider  $p = 0$  in (2.6), then we get the following classical Pochhammer symbol defined by

$$(\delta)_n = \frac{\Gamma(\delta+n)}{\Gamma(\delta)} = \begin{cases} \delta(\delta+1)(\delta+2)\cdots(\delta+n-1), & (n \in \mathbb{N}, \delta \in \mathbb{C}), \\ 1, & (n = 0, \delta \in \mathbb{C} \setminus 0), \end{cases}$$

where  $\Gamma(\delta)$  is defined by

$$\Gamma(\delta) = \int_0^{\infty} t^{\delta-1} e^{-t} dt.$$

The extensions and applications of these Mittag-Leffler functions were extensively studied by the researchers ([19, 28, 37–41, 43]).

The Lebesgue measurable space for real (complex) valued functions is defined by

$$\mathcal{L}(x_1, x_2) = \left\{ h : \|h\|_1 = \int_{x_1}^{x_2} |h(x)| dx < \infty \right\}. \quad (2.7)$$

The Riemann-Liouville type (left and right sided) fractional integral operators are respectively defined in [11, 35] by

$$(\mathfrak{I}_{x_1+}^{\kappa} h)(x) = \frac{1}{\Gamma(\kappa)} \int_{x_1}^x \frac{h(v)}{(x-v)^{1-\kappa}} dv, \quad (x > x_1), \quad (2.8)$$

and

$$(\mathfrak{I}_{x_2^-}^\kappa h)(x) = \frac{1}{\Gamma(\kappa)} \int_x^{x_2} \frac{h(v)}{(v-x)^{1-\kappa}} dv, \quad (x < x_2),$$

where  $h \in L(x_1, x_2)$ ,  $\kappa \in \mathbb{C}$  and  $\Re(\kappa) > 0$ .

The Riemann-Liouville fractional derivatives (left and right sided) for  $h(x) \in L(x_1, x_2)$ ,  $\kappa \in \mathbb{C}$ ,  $\Re(\kappa) > 0$  and  $n = [\Re(\kappa)] + 1$  are defined in [11, 35] by

$$(\mathfrak{D}_{x_1^+}^\kappa h)(x) = \left(\frac{d}{dx}\right)^n (\mathfrak{I}_{x_1^+}^{n-\kappa} h)(x), \quad (2.9)$$

$$(\mathfrak{D}_{x_2^-}^\kappa h)(x) = \left(-\frac{d}{dx}\right)^n (\mathfrak{I}_{x_2^-}^{n-\kappa} h)(x),$$

respectively. The generalized differential operator  $\mathfrak{D}_{x_1^+}^{\kappa, \nu}$  of order  $0 < \kappa < 1$  and type  $0 < \nu < 1$  with respect to  $x$  can be found in ([25, 35]) as;

$$(\mathfrak{D}_{x_1^+}^{\kappa, \nu} h) = \left( \mathfrak{I}_{x_1^+}^{\nu(1-\lambda)} \frac{d}{dx} (\mathfrak{I}_{x_1^+}^{(1-\nu)(1-\lambda)} h) \right) (x). \quad (2.10)$$

Clearly, if we put  $\nu = 0$  then (2.10) reduces to the operator  $\mathfrak{D}_{x_1^+}^\kappa$  defined in (2.9).

Moreover, we consider the following well known results.

**Theorem 2.1.** [26] *The following results holds true:*

$$\mathfrak{I}_{x_1^+}^\kappa (\nu - x_1)^{\varphi-1} = \frac{\Gamma(\varphi)}{\Gamma(\kappa + \varphi)} (x - x_1)^{\kappa+\varphi-1}, \quad (2.11)$$

where  $\kappa, \varphi \in \mathbb{C}$ ,  $\Re(\kappa) > 0$ ,  $\Re(\varphi) > 0$ ,

**Theorem 2.2.** [42] *Suppose that the function  $h(z)$  has a power series expansion  $h(z) = \sum_{k=0}^{\infty} k_n z^k$  and analytic in the disc  $|z| < R$ , then the following result holds:*

$${}_0\mathfrak{D}_z^\kappa \{z^{\varphi-1} h(z)\} = \frac{\Gamma(\varphi)}{\Gamma(\kappa + \varphi)} \sum_{n=0}^{\infty} \frac{a_n(\varphi)_n}{(\kappa + \varphi)_n} z^n.$$

**Lemma 2.1.** [43] *Suppose that  $x > x_1$ ,  $\kappa \in (0, 1)$ ,  $\nu \in [0, 1]$ ,  $\Re(\varphi) > 0$  and  $\Re(\kappa) > 0$ , then we have*

$$\mathfrak{D}_{x_1^+}^\kappa [(\nu - x_1)^{\varphi-1}](x) = \frac{\Gamma(\varphi)}{\Gamma(\varphi - \kappa)} (x - x_1)^{\varphi-\kappa-1}. \quad (2.12)$$

### 3. Fractional integral and differential formulas of an extended Mittag-Leffler function

In this section, we present some fractional integration and differentiation formulas of an extended Mittag-Leffler function (2.4). We consider the following well-known result recently presented by Rahman et al. [29].

**Theorem 3.1.** *The following differentiation holds for  $\delta, \kappa, \varphi \in \mathbb{C}$ ,  $\Re(\varphi) > 0$ ,  $\Re(\kappa) > 0$  and  $n \in \mathbb{N}$ ,*

$$\left(\frac{d}{dz}\right)^n \left(z^{\kappa-1} E_{\rho, \varphi; p, \nu}^\delta(\omega z^\rho)\right) = z^{\kappa-n-1} E_{\rho, \varphi-n; p, \nu}^\delta(\omega z^\rho). \quad (3.1)$$

Next, we present the following fractional calculus approach.

**Theorem 3.2.** Suppose  $x > x_1 (x_1 \in \mathbb{R}_+ = [0, \infty))$ ,  $\delta, \kappa, \varphi, \omega \in \mathbb{C}$ ,  $\Re(\varphi) > 0$  and  $\Re(\kappa) > 0$ , then

$$\mathfrak{I}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})](x) = (x - x_1)^{\kappa+\varphi-1} E_{\rho, \varphi+\kappa; p, v}^{\delta}(\omega(x - x_1)^{\rho}), \quad (3.2)$$

$$\mathfrak{D}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(x - x_1)^{\rho})] = (x - x_1)^{\varphi-\kappa-1} E_{\rho, \varphi-\kappa; p, v}^{\delta}(\omega(x - x_1)^{\rho}) \quad (3.3)$$

and

$$\mathfrak{D}_{x_1+}^{\kappa, v}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(x - x_1)^{\rho})](x) = (x - x_1)^{\varphi-\kappa-1} E_{\rho, \varphi-\kappa; p, v}^{\delta}(\omega(x - x_1)^{\rho}). \quad (3.4)$$

*Proof.* Consider

$$\begin{aligned} & \mathfrak{I}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(x - x_1)^{\rho})] \\ &= \frac{1}{\Gamma(\kappa)} \int_{x_1}^x \frac{(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})}{(x - v)^{1-\kappa}} dv \\ &= \frac{1}{\Gamma(\kappa)} \sum_{n=0}^{\infty} \frac{(\delta; p, v)_n \omega^n}{\Gamma(\rho n + \varphi) n!} \int_{x_1}^x (v - x_1)^{\varphi+\rho n-1} (x - v)^{\kappa-1} dv \\ &= \sum_{n=0}^{\infty} \frac{(\delta; p, v)_n \omega^n}{\Gamma(\rho n + \varphi) n!} \left( \mathfrak{I}_{x_1+}^{\kappa}[(v - x_1)^{\varphi+\rho n-1}] \right). \end{aligned}$$

By the use of (2.11), we have

$$\begin{aligned} & \mathfrak{I}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(x - x_1)^{\rho})] \\ &= \sum_{n=0}^{\infty} \frac{(\delta; p, v)_n \omega^n}{\Gamma(\rho n + \varphi) n!} (x - x_1)^{\varphi+\kappa+\rho n-1} \cdot \frac{\Gamma(\rho n + \varphi)}{\Gamma(\rho n + \varphi + \kappa)} \\ &= (x - x_1)^{\varphi+\kappa-1} \sum_{n=0}^{\infty} \frac{(\delta; p, v)_n}{\Gamma(\rho n + \varphi + \kappa)} \frac{[\omega(x - x_1)^{\rho}]^n}{n!} \\ &= (x - x_1)^{\varphi+\kappa-1} E_{\rho, \varphi+\kappa; p, v}^{\delta}(\omega(x - x_1)^{\rho}), \end{aligned}$$

which completes the proof of (3.2).

Now, we have

$$\mathfrak{D}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})] = \left(\frac{d}{dx}\right)^n \left\{ \mathfrak{I}_{x_1+}^{n-\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})] \right\},$$

which on using (3.2) takes the following form:

$$\mathfrak{D}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})] = \left(\frac{d}{dx}\right)^n \left\{ (x - x_1)^{\varphi-\kappa+n-1} E_{\rho, \varphi-\kappa+n; p, v}^{\delta}(\omega(x - x_1)^{\rho}) \right\}.$$

Applying (3.1), we have

$$\mathfrak{D}_{x_1+}^{\kappa}[(v - x_1)^{\varphi-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(v - x_1)^{\rho})](x) = \left\{ (x - x_1)^{\varphi-\kappa-1} E_{\rho, \varphi-\kappa; p, v}^{\delta}(\omega(v)^{\rho}) \right\}.$$

This completes the proof of (3.3).

To prove (3.4), we have

$$\begin{aligned} & \left( \mathfrak{D}_{x_1+}^{\kappa,v}[(v-x_1)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(v-x_1)^{\rho})] \right)(x) \\ &= \left( \mathfrak{D}_{x_1+}^{\kappa,v} \left[ \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{\omega^n}{n!} (v-x_1)^{\rho n + \varphi - 1} \right] \right)(x) \\ &= \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{\omega^n}{n!} \left( \mathfrak{D}_{x_1+}^{\kappa,v}[(v-x_1)^{\rho n + \varphi - 1}] \right)(x). \end{aligned}$$

By applying (2.12), we get

$$\begin{aligned} & \left( \mathfrak{D}_{x_1+}^{\kappa,v}[(v-x_1)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(v-x_1)^{\rho})] \right)(x) \\ &= \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{\omega^n}{n!} \frac{\Gamma(\rho n + \varphi)}{\Gamma(\rho n + \varphi - \kappa)} (x-x_1)^{\rho n + \varphi - \kappa - 1} \\ &= (x-x_1)^{\varphi - \kappa - 1} \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi - \kappa)} \frac{[\omega(x-x_1)^{\rho}]^n}{n!} \\ &= (x-x_1)^{\varphi - \kappa - 1} E_{\rho,\varphi-\kappa;p,v}^{\delta}(\omega(x-x_1)^{\rho}), \end{aligned}$$

which completes the required proof.  $\square$

If we consider  $v = 0$ , then we get the following result of Parmar et al. [30].

**Corollary 3.1.** Suppose  $x > x_1$  ( $x_1 \in \mathbb{R}_+ = [0, \infty)$ ),  $\delta, \kappa, \varphi, \omega \in \mathbb{C}$ ,  $\Re(\varphi) > 0$  and  $\Re(\kappa) > 0$ , then

$$\mathfrak{I}_{x_1+}^{\kappa}[(v-x_1)^{\varphi-1} E_{\rho,\varphi;p}^{\delta}(\omega(v-x_1)^{\rho})](x) = (x-x_1)^{\kappa+\varphi-1} E_{\rho,\varphi+\kappa;p}^{\delta}(\omega(x-x_1)^{\rho}),$$

$$\mathfrak{D}_{x_1+}^{\kappa}[(v-x_1)^{\varphi-1} E_{\rho,\varphi;p}^{\delta}(\omega(x-x_1)^{\rho})] = (x-x_1)^{\varphi-\kappa-1} E_{\rho,\varphi-\kappa;p}^{\delta}(\omega(x-x_1)^{\rho})$$

and

$$\mathfrak{D}_{x_1+}^{\kappa,v}[(v-x_1)^{\varphi-1} E_{\rho,\varphi;p}^{\delta}(\omega(x-x_1)^{\rho})](x) = (x-x_1)^{\varphi-\kappa-1} E_{\rho,\varphi-\kappa;p}^{\delta}(\omega(x-x_1)^{\rho}).$$

**Remark 3.1.** Put  $p = 0$  in Corollary 3.1, then we lead to the work presented by Prabhakar [31].

#### 4. Extension of Prabhakar type fractional integral and its properties

In this section, we estimate a new extension of Prabhakar type fractional integral and its properties.

**Definition 4.1.** Let  $\varphi, \rho, \delta, \omega \in \mathbb{C}$ ,  $\Re(\delta) > 0$ ,  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ . Then the new Prabhakar left and right sided fractional integral are defined by

$$({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h)(x) = \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) h(v) dv, \quad x > x_1 \quad (4.1)$$

and

$$({}_p^v \mathcal{E}_{x_2-;\rho,\varphi}^{\omega;\delta} h)(x) = \int_x^{x_2} (v-x)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(v-x_2)^{\rho}) h(v) dv, \quad x < x_2, \quad (4.2)$$

respectively. Substituting  $v = 0$ , then (4.1) and (4.2) reduce to the following operators

$$({}_p\mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta}h)(x) = \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi;p}^{\delta}(\omega(x-v)^{\rho})h(v)dv, x > x_1 \quad (4.3)$$

and

$$({}_p\mathcal{E}_{x_2-;\rho,\varphi}^{\omega;\delta}h)(x) = \int_x^{x_2} (v-x)^{\varphi-1} E_{\rho,\varphi;p}^{\delta}(\omega(v-x_2)^{\rho})h(v)dv, x < x_2. \quad (4.4)$$

When we substitute  $p = 0$ , then (4.3) and (4.4) reduce to the following Prabhakar operators

$$(\mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta}h)(x) = \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi}^{\delta}(\omega(x-v)^{\rho})h(v)dv, x > x_1 \quad (4.5)$$

and

$$(\mathcal{E}_{x_2-;\rho,\varphi}^{\omega;\delta}h)(x) = \int_x^{x_2} (v-x)^{\varphi-1} E_{\rho,\varphi}^{\delta}(\omega(v-x_2)^{\rho})h(v)dv, x < x_2. \quad (4.6)$$

If we consider  $\omega = 0$ , then the integral operators in (4.5) and (4.6) reduce to the well-known Riemann-Liouville fractional integral operators.

Next, we present the following results.

**Theorem 4.1.** Suppose that  $\varphi, \rho, \kappa, \delta, \omega \in \mathbb{C}$ ,  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ ,  $\Re(\kappa) > 0$ ,  $p, v \geq 0$  and  $\Re(\delta) > 0$ , then the following result holds:

$$\left( {}_p^v\mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta}[(v-x_1)^{\kappa-1}] \right)(x) = (x-x_1)^{\kappa+\varphi-1} \Gamma(\kappa) E_{\rho,\varphi+\kappa}^{\delta;p,v}(\omega(x-x_1)^{\rho}).$$

*Proof.* From definition of Prabhakar fractional integral (4.1)

$$\left( {}_p^v\mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta}h \right)(x) = \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi}^{\delta}(\omega(x-v)^{\rho})h(v)dv.$$

Therefore, we have

$$\begin{aligned} \left( {}_p^v\mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta}[(v-x_1)^{\kappa-1}] \right)(x) &= \int_{x_1}^x (x-v)^{\varphi-1} (v-x_1)^{\kappa-1} E_{\rho,\varphi}^{\delta;p,v}(\omega(x-v)^{\rho})dv \\ &= \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{\omega^n}{n!} \left( \int_{x_1}^x (v-x_1)^{\kappa-1} (x-v)^{\kappa+\rho n-1} dv \right) \\ &= \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{\omega^n}{n!} \mathfrak{I}_{x_1+}^{\rho n + \varphi} [(v-x_1)^{\kappa-1}] \\ &= (x-x_1)^{\varphi+\kappa-1} \left\{ \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n}{\Gamma(\rho n + \varphi)} \frac{[\omega(x-x_1)^{\rho}]^n}{n!} \frac{\Gamma(\kappa)\Gamma(\rho n + \varphi)}{\Gamma(\rho n + \varphi + \kappa)} \right\} \\ &= (x-x_1)^{\varphi+\kappa-1} \Gamma(\kappa) E_{\rho,\varphi+\kappa}^{\delta;p,v}(\omega(x-x_1)^{\rho}), \end{aligned}$$

which completes the desired proof.  $\square$

**Theorem 4.2.** Suppose that  $\delta, \rho, \varphi, \omega \in \mathbb{C}$ ,  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ ,  $v, p \geq 0$ , then the following result holds:

$$\| {}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} \Phi \|_1 \leq M \| \Phi \|_1,$$

where

$$M = (x_2 - x_1)^{\Re(\varphi)} \sum_{n=0}^{\infty} \frac{|(\delta; p, v)_n|}{\Gamma(\rho n + \varphi)(\Re(\varphi) + \Re(\rho)n)} \frac{|\omega(x_2 - x_1)^\rho|^n}{n!}.$$

*Proof.* Consider (4.1) and (2.7) and interchanging the order of integrations, we obtain

$$\begin{aligned} \| {}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} \Phi \|_1 &= \int_{x_1}^{x_2} \left| \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho, \varphi}^{\delta}(\omega(x-v)^\rho) \Phi(v) dv \right| dx \\ &\leq \int_{x_1}^{x_2} \left[ \int_v^x (x-v)^{\Re(\varphi)-1} |E_{\rho, \varphi; p, v}^{\delta}(\omega(x-v)^\rho)| dx \right] |\Phi(v)| dv \\ &= \int_{x_1}^{x_2} \left[ \int_0^{x-v} (u)^{\Re(\varphi)-1} |E_{\rho, \varphi; p, v}^{\delta}(\omega(u)^\rho)| du \right] |\Phi(v)| dv, \text{ (By setting } (x-v) = u \text{).} \end{aligned}$$

After simplification, we obtain

$$\| {}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} \Phi \|_1 \leq \int_{x_1}^{x_2} \left[ \sum_{n=0}^{\infty} \frac{|(\delta; p, v)_n| |\omega^n|}{\Gamma(\rho n + \varphi) n!} \left( \frac{(u)^{\Re(\varphi) + \Re(\rho)n}}{(\Re(\varphi) + \Re(\rho)n)} \right)_0^{x_2 - x_1} \right] |\Phi(v)| dv.$$

This can further be expressed as

$$\begin{aligned} \| {}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} \Phi \|_1 &\leq \left\{ (x_2 - x_1)^{\Re(\varphi)} \sum_{n=0}^{\infty} \frac{|(\delta; p, v)_n|}{\Gamma(\rho n + \varphi)(\Re(\varphi) + \Re(\rho)n)} \frac{|\omega(x_2 - x_1)^\rho|^n}{n!} \right\} \cdot \int_{x_1}^{x_2} |\Phi(v)| dv \\ &= M \| \Phi \|_1, \end{aligned}$$

where

$$M = (x_2 - x_1)^{\Re(\varphi)} \sum_{n=0}^{\infty} \frac{|(\delta; p, v)_n|}{\Gamma(\rho n + \varphi)(\Re(\varphi) + \Re(\rho)n)} \frac{|\omega(x_2 - x_1)^\rho|^n}{n!},$$

which completes the desired result.  $\square$

**Corollary 4.1.** The following result holds for  $\rho, \varphi, \delta \in \mathbb{C}$ ,  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ ,  $\Re(\delta) > 0$ ,

$$\left( {}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} 1 \right)(x) = (x - x_1)^\varphi E_{\rho, \varphi+1; p, v}^{\delta}(\omega(x - x_1)^\rho).$$

*Proof.* Consider



$$\begin{aligned}({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} 1)(x) &= \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) dv \\ &= \int_{x_1}^x (x-v)^{\varphi-1} \left( \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n \omega^n (x-v)^{n\rho}}{\Gamma(n\rho+\varphi)n!} \right) dv.\end{aligned}$$

Interchanging the order of summation and integration, we have

$$\begin{aligned}({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} 1)(x) &= \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n \omega^n}{\Gamma(n\rho+\varphi)n!} \int_{x_1}^x (x-v)^{\varphi+n\rho-1} dv \\ &= (x-x_1)^{\varphi} \sum_{n=0}^{\infty} \frac{(\delta;p,v)_n \omega^n (x-x_1)^{n\rho}}{\Gamma(n\rho+\varphi)(n\rho+\varphi)n!} \\ &= (x-x_1)^{\varphi} E_{\rho,\varphi+1;p,v}^{\delta}(\omega(x-x_1)^{\rho}),\end{aligned}$$

which completes the desired proof.  $\square$

**Theorem 4.3.** *The extended Prabhakar operator can be expressed in the following series of Riemann-Liouville fractional integrals for  $\delta, \rho, \varphi, \omega \in \mathbb{C}$  with  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ ,  $\Re(\delta) > 0$ ,  $p, v \geq 0$  and  $x > x_1$ ,*

$$({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h)(x) = \sum_{n=0}^{\infty} \frac{\Gamma_v(\delta+n;p)\omega^n}{\Gamma(\delta)} \mathfrak{I}_{x_1+}^{\rho n+\varphi} h(x).$$

*Proof.* By using (2.4) in (4.1) and then interchanging the order of summation and integration, we obtain

$$\begin{aligned}({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h)(x) &= \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) h(v) dv \\ &= \int_{x_1}^x (x-v)^{\varphi-1} \sum_{n=0}^{\infty} \frac{\Gamma_v(\delta+n;p)\omega^n (x-v)^{n\rho}}{\Gamma(\rho n+\varphi)} h(v) dv \\ &= \sum_{n=0}^{\infty} \frac{\Gamma_v(\delta+n;p)\omega^n}{\Gamma(\delta)} \frac{1}{\Gamma(\rho n+\varphi)} \int_{x_1}^x (x-v)^{\rho n+\varphi-1} h(v) dv \\ &= \sum_{n=0}^{\infty} \frac{\Gamma_v(\delta+n;p)\omega^n}{\Gamma(\delta)} \mathfrak{I}_{x_1+}^{\rho n+\varphi} h(x),\end{aligned}$$

which completes the desired proof.  $\square$

**Theorem 4.4.** *The following result satisfies for  $\kappa, \delta, \rho, \varphi, \omega \in \mathbb{C}$  with  $\Re(\rho) > 0$ ,  $\Re(\varphi) > 0$ ,  $\Re(\delta) > 0$ ,  $\Re(\kappa) > 0$ ,  $p, v \geq 0$  and  $x > x_1$ ,*

$$(\mathfrak{I}_{x_1+}^{\kappa} [{}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) = ({}_p^v \mathcal{E}_{x_1+;\rho,\varphi+\kappa}^{\omega;\delta} h)(x) = ({}_p^v \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} [\mathfrak{I}_{x_1+}^{\kappa} h])(x), \quad (4.7)$$

for arbitrary function  $h \in L(x_1, x_2)$ .

*Proof.* By utilizing (4.1) and (2.8), we have

$$\begin{aligned} (\mathfrak{I}_{x_1+}^{\kappa} [{}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) &= \frac{1}{\Gamma(\kappa)} \int_{x_1}^x \frac{[({}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h)(v)]}{(x-v)^{1-\kappa}} dv \\ &= \frac{1}{\Gamma(\kappa)} \int_{x_1}^x (x-v)^{\kappa-1} \left[ \int_{x_1}^v (v-u)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(v-u)^{\rho}) h(u) du \right] dv. \end{aligned}$$

By interchanging the order of integrations, we have

$$(\mathfrak{I}_{x_1+}^{\kappa} [{}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) = \int_{x_1}^x \left[ \frac{1}{\Gamma(\kappa)} \int_u^x (x-v)^{\kappa-1} (v-u)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(v-u)^{\rho}) dv \right] h(u) du.$$

By putting  $v-u = \theta$ , we obtain

$$\begin{aligned} (\mathfrak{I}_{x_1+}^{\kappa} [{}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) &= \int_{x_1}^x \left[ \frac{1}{\Gamma(\kappa)} \int_0^{x-u} (x-u-\theta)^{\kappa-1} (\theta)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(\theta)^{\rho}) d\theta \right] h(u) du \\ &= \int_{x_1}^x \left[ \frac{1}{\Gamma(\kappa)} \int_0^{x-u} \frac{(\theta)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(\theta)^{\rho})}{(x-u-\theta)^{1-\kappa}} d\theta \right] h(u) du. \end{aligned}$$

Thus by the use of (2.8) and applying (3.2), we obtain

$$\begin{aligned} (\mathfrak{I}_{x_1+}^{\kappa} [{}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) &= \int_{x_1}^x \left[ (\theta)^{\kappa+\varphi-1} E_{\rho,\varphi+\kappa;p,v}^{\delta}(\omega(\theta)^{\rho}) \right] h(u) du \\ &= \int_{x_1}^x (x-u)^{\kappa+\varphi-1} E_{\rho,\varphi+\kappa}^{\delta}(\omega(x-u)^{\rho}) h(u) du. \end{aligned}$$

Thus, we obtain

$$(\mathfrak{I}_{x_1+}^{\kappa} [{}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} h])(x) = ({}^v_p \mathcal{E}_{x_1+;\rho,\varphi+\kappa}^{\omega;\delta} h)(x). \quad (4.8)$$

Next, consider the right hand side of (4.7) and utilizing (4.1) to derive the second part, we have

$$\begin{aligned} ({}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} [\mathfrak{I}_{x_1+}^{\kappa} h])(x) &= \int_{x_1}^x (x-v)^{\varphi-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) [\mathfrak{I}_{x_1+}^{\kappa} h](v) dv \\ &= \int_{x_1}^x E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) \left( \frac{1}{\Gamma(\kappa)} \int_{x_1}^v \frac{f(u)}{(v-u)^{1-\kappa}} du \right) dv. \end{aligned}$$

By interchanging the order of integrations, we obtain

$$({}^v_p \mathcal{E}_{x_1+;\rho,\varphi}^{\omega;\delta} [\mathfrak{I}_{x_1+}^{\kappa} h])(x) = \int_{x_1}^x \frac{1}{\Gamma(\kappa)} \left[ \int_u^x (x-v)^{\varphi-1} (v-u)^{\kappa-1} E_{\rho,\varphi;p,v}^{\delta}(\omega(x-v)^{\rho}) dv \right] h(u) du.$$

By setting  $x - \tau = \theta$ , we have

$$\begin{aligned}({}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} [\mathfrak{I}_{x_1+}^{\kappa} h])(x) &= \int_{x_1}^x \frac{1}{\Gamma(\kappa)} \left[ \int_{x-u}^0 (\theta)^{\varphi-1} (x - \theta - u)^{\kappa-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(\theta)^{\rho})(-d\theta) \right] h(u) du \\ &= \int_a^x \frac{1}{\Gamma(\kappa)} \left[ \int_0^{x-u} (\theta)^{\varphi-1} (x - \theta - u)^{\kappa-1} E_{\rho, \varphi; p, v}^{\delta}(\omega(\theta)^{\rho}) d\theta \right] h(u) du.\end{aligned}$$

Further, by using (2.8) and applying (3.2), we obtain

$$({}_p^v \mathcal{E}_{x_1+; \rho, \varphi}^{\omega; \delta} [\mathfrak{I}_{x_1+}^{\kappa} h])(x) = ({}_p^v \mathcal{E}_{x_1+; \rho, \varphi + \kappa}^{\omega; \delta} h)(x). \quad (4.9)$$

Thus (4.8) and (4.9) complete the desired proof.  $\square$

## 5. Conclusions

In this article, we have investigated fractional integral and differential formulas for the extended Mittag-Leffler function. Also, we defined a further extension of Prabhakar fractional operator and investigated its various properties. The special cases of our result can be found in the literature cited therein. One can easily investigate many of its applications and develop various fractional integral inequalities by utilizing the newly defined Prabhakar fractional operator.

## Authors contributions

All the authors contributed equally and they read and approved the final manuscript for publication.

## Conflict of interest

The authors declare that they have no competing interests.

## References

1. T. Abdeljawad, On conformable fractional calculus, *J. Comput. Appl. Math.*, **279** (2015), 57–66.
2. T. Abdeljawad, D. Baleanu, Monotonicity results for fractional difference operators with discrete exponential kernels, *Adv. Differ. Equations*, **2017** (2017), 78.
3. T. Abdeljawad, D. Baleanu, On fractional derivatives with exponential kernel and their discrete versions, *Rep. Math. Phys.*, **80** (2017), 11–27.
4. D. R. Anderson, D. J. Ulness, Newly defined conformable derivatives, *Adv. Dyn. Syst. Appl.*, **10** (2015), 109–137.
5. A. Atangana, D. Baleanu, New fractional derivatives with nonlocal and non-singular kernel: Theory and application to heat transfer model, *Therm. Sci.*, **20** (2016), 763–769.
6. M. Caputo, M. Fabrizio, A new definition of fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.*, **1** (2015), 1–13.

7. F. Jarad, T. Abdeljawad, J. Alzabut, Generalized fractional derivatives generated by a class of local proportional derivatives, *Eur. Phys. J. Spec. Top.*, **226** (2017), 3457–3471.
8. F. Jarad, E. Uğurlu, T. Abdeljawad, D. Baleanu, On a new class of fractional operators, *Adv. Differ. Equations*, **2017** (2017), 247.
9. R. Khalil, M. Al Horani, A. Yousef, M. Sababheh, A new definition of fractional derivative, *J. Comput. Appl. Math.*, **264** (2014), 65–70.
10. J. Losada, J. J. Nieto, Properties of a new fractional derivative without singular kernel, *Progr. Fract. Differ. Appl.*, **1** (2015), 87–92.
11. I. Podlubny, *Fractional differential equations*, London: Academic Press, 1999.
12. R. Hilfer, *Applications of fractional calculus in physics*, Singapore: World Scientific, 2000.
13. D. Baleanu, K. Diethelm, E. Scalas, J. J. Trujillo, *Fractional calculus models and numerical methods*, Singapore: World Scientific, 2012.
14. D. Baleanu, D. Kumar, S. D. Purohit, Generalized fractional integrals of product of two  $H$ -functions and a general class of polynomials, *Int. J. Comput. Math.*, **93** (2016), 1320–1329.
15. R. F. Camargo, E. C. de Oliveira, J. Vas, On the generalized Mittag-Leffler function and its application in a fractional telegraph equation, *Math. Phys. Anal. Geom.*, **15** (2012), 1–16.
16. M. A. Chaudhry, S. M. Zubair, *On a class of incomplete gamma functions with applications*, Boca Raton: Chapman and Hall, 2001.
17. M. M. Džrbašjan, *Integral transforms and representations of functions in the complex domain*, Moscow: Nauka, 1966.
18. J. F. Gómez-Aguilar, A. Atangana, Fractional Hunter-Saxton equation involving partial operators with bi-order in Riemann-Liouville and Liouville-Caputo sense, *Eur. Phys. J. Plus*, **132** (2017), 100.
19. R. Gorenflo, A. A. Kilbas, S. V. Rogosin, On the generalized Mittag-Leffler type functions, *Integr. Transf. Spec. Funct.*, **7** (1998), 215–224.
20. R. Gorenflo, Y. Luchko, F. Mainardi, Wright functions as scale-invariant solutions of the diffusion-wave equation, *J. Comput. Appl. Math.*, **118** (2000), 175–191.
21. R. Gorenflo, F. Mainardi, Integral and differential equations of fractional order, *Fractals Fractional Calculus Continuum Mech.*, (2000), 223–276.
22. R. Gorenflo, F. Mainardi, H. M. Srivastava, Special functions in fractional relaxation oscillation and fractional diffusion-wave phenomena, In: D. Bainov, *Proceedings of the eighth international colloquium on differential equations*, (1998), 195–202.
23. R. Hilfer, H. Seybold, Computation of the generalized Mittag-Leffler function and its inverse in the complex plane, *Integr. Transf. Spec. Funct.*, **17** (2006), 637–652.
24. A. A. Kilbas, M. Saigo, On Mittag-Leffler type function, fractional calculus operators and solutions of integral equations, *Integr. Transf. Spec. Funct.*, **4** (1996), 355–370.
25. A. A. Kilbas, H. M. Srivastava, J. J. Trujillo, *Theory and applications of fractional differential equations*, Amsterdam: Elsevier Science B.V., 2006.

26. A. M. Mathai, H. J. Haubold, *Special functions for applied scientists*, New York: Springer, 2008.
27. K. S. Nisar, S. D. Purohit, S. R. Mondal, Generalized fractional kinetic equations involving generalized Struve function of the first kind, *J. King Saud Univ.-Sci.*, **28** (2016), 167–171.
28. M. A. Özarslan, B. Yılmaz, The extended Mittag-Leffler function and its properties, *J. Inequa. Appl.*, **2014** (2014), 85.
29. G. Rahman, A. Saboor, Z. Anjum, K. S. Nisar, T. Abdeljawad, An extension of the Mittag-Leffler function and its associated properties, *Adv. Math. Phys.*, **2020** (2020), 5792853.
30. J. Choi, R. K. Parmar, P. Chopra, Extended Mittag-Leffler function and associated fractional calculus operators, *Georgian Math. J.*, **27** (2019), 1–11.
31. T. R. Prabhakar, A singular integral equation with a generalized Mittag-Leffler function in the kernel, *Yokohama Math. J.*, **19** (1971), 7–15.
32. S. D. Purohit, Solutions of fractional partial differential equations of quantum mechanics, *Adv. Appl. Math. Mech.*, **5** (2013), 639–651.
33. S. D. Purohit, S. L. Kalla, On fractional partial differential equations related to quantum mechanics, *J. Phys. A: Math. Theor.*, **44** (2011), 045202.
34. M. Saigo, A. A. Kilbas, On Mittag-Leffler type function and applications, *Integr. Transf. Spec. Funct.*, **7** (1998), 97–112.
35. S. G. Samko, A. A. Kilbas, O. I. Marichev, *Fractional integrals and derivatives: Theory and applications*, Switzerland: Gordon and Breach, 1993.
36. H. J. Seybold, R. Hilfer, Numerical results for the generalized Mittag-Leffler function, *Fract. Calc. Appl. Anal.*, **8** (2005), 127–139.
37. A. K. Shukla, J. C. Prajapati, On a generalization of Mittag-Leffler functions and its properties, *J. Math. Anal. Appl.*, **336** (2007), 797–811.
38. H. M. Srivastava, A contour integral involving Fox's  $H$ -function, *Indian J. Math.*, **14** (1972), 1–6.
39. H. M. Srivastava, A note on the integral representation for the product of two generalized Rice polynomials, *Collect. Math.*, **24** (1973), 117–121.
40. H. M. Srivastava, K. C. Gupta, S. P. Goyal, *The  $H$ -functions of one and two variables with applications*, New Delhi: South Asian Publishers, 1982.
41. H. M. Srivastava, C. M. Joshi, Integral representation for the product of a class of generalized hypergeometric polynomials, *Bull. Acad. R. Belg.*, **60** (1974), 919–926.
42. H. M. Srivastava, H. L. Manocha, *A treatise on generating functions*, New York: Halsted Press, 1984.
43. H. M. Srivastava, Ž. Tomovski, Fractional calculus with an integral operator containing a generalized Mittag-Leffler function in the kernel, *Appl. Math. Comput.*, **211** (2009), 189–210.
44. H. M. Srivastava, G. Rahman, K. S. Nisar, Some extensions of the Pochhammer symbol and the associated hypergeometric functions, *Iran. J. Sci. Technol. Trans. Sci.*, **43** (2019), 2601–2606.
45. H. M. Srivastava, A. Çetinkaya, İ. Onur Kıymaz, A certain generalized Pochhammer symbol and its applications to hypergeometric functions, *Appl. Math. Comput.*, **226** (2014), 484–491.

46. M. Z. Sarikaya, H. Budak, F. Usta, On generalized the conformable fractional calculus, *TWMS J. Appl. Eng. Math.*, **9** (2019), 792–799.
47. F. Usta, H. Budak, M. Z. Sarikaya, Some new Chebyshev type inequalities utilizing generalized fractional integral operators, *AIMS Math.*, **5** (2020), 1147–1161.
48. F. Usta, M. Z. Sarikaya, On generalization conformable fractional integral inequalities, *Filomat*, **32** (2018), 5519–5526.
49. M. Z. Sarikaya, F. Usta, On comparison theorems for conformable fractional differential equations, *Int. J. Anal. Appl.*, **12** (2016), 207–214.
50. F. Usta, M. Z. Sarikaya, F. Qi, Explicit bounds on certain integral inequalities via conformable fractional calculus, *Cogent Math.*, **4** (2017), 1277505.
51. F. Usta, M. Z. Sarikaya, Some improvements of conformable fractional integral inequalities, *Int. J. Anal. Appl.*, **14** (2017), 162–166.
52. F. Usta, M. Z. Sarikaya, On bivariate retarded integral inequalities and their applications, *Facta Univ.-Ser.: Math. Inf.*, **34** (2019), 553–561.
53. G. M. Mittag-Leffler, Sur la nouvelle fonction  $E_\alpha(z)$ , *C. R. Acad. Sci. Paris*, **137** (1903), 554–558.
54. A. Wiman, Über den fundamental satz in der theorie der functionen  $E_\alpha(z)$ , *Acta Math.*, **29** (1905), 191–201.



AIMS Press

©2021 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)