



---

*Research article*

## Quantum Hermite-Hadamard and quantum Ostrowski type inequalities for $s$ -convex functions in the second sense with applications

Suphawat Asawasamrit<sup>1</sup>, Muhammad Aamir Ali<sup>2,\*</sup>, Hüseyin Budak<sup>3</sup>, Sotiris K. Ntouyas<sup>4,5</sup> and Jessada Tariboon<sup>1,\*</sup>

<sup>1</sup> Intelligent and Nonlinear Dynamic Innovations Research Center, Department of Mathematics, Faculty of Applied Sciences, King Mongkut's University of Technology North Bangkok, Bangkok 10800, Thailand

<sup>2</sup> Jiangsu Key Laboratory for NSLSCS, School of Mathematical Sciences, Nanjing Normal University, Nanjing 210023, China

<sup>3</sup> Department of Mathematics, Faculty of Arts and Sciences, Düzce University, Düzce, Turkey

<sup>4</sup> Department of Mathematics, University of Ioannina, 45110 Ioannina, Greece

<sup>5</sup> Nonlinear Analysis and Applied Mathematics (NAAM)-Research Group, Department of Mathematics, Faculty of Sciences, King Abdulaziz University, P. O. Box 80203, Jeddah 21589, Saudi Arabia

\* **Correspondence:** Email: mahr.muhammad.aamir@gmail.com, jessada.t@sci.kmutnb.ac.th.

**Abstract:** In this study, we use quantum calculus to prove Hermite-Hadamard and Ostrowski type inequalities for  $s$ -convex functions in the second sense. The newly proven results are also shown to be an extension of comparable results in the existing literature. Furthermore, it is provided that how the newly discovered inequalities can be applied to special means of real numbers.

**Keywords:** Hermite-Hadamard inequality; Ostrowski inequality;  $q$ -integral; quantum calculus;  $s$ -convex functions

**Mathematics Subject Classification:** 26D10, 26A51, 26D15

---

## 1. Introduction

In convex functions theory, Hermite-Hadamard (H-H) inequality is very important and was discovered by C. Hermite and J. Hadamard independently (see, also [23], [41, p.137]),

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x)dx \leq \frac{f(a)+f(b)}{2} \quad (1.1)$$

where  $f$  is a convex function. In the case of concave mappings, the above inequality is satisfied in reverse order. For more recent developments, one can consult [17, 18, 42, 48, 49].

Hudzik and Maligranda defined  $s$ -convex functions in the second sense in [28], which may be expressed as: a mapping  $f : \mathbb{R}^+ \rightarrow \mathbb{R}$ , where  $\mathbb{R}^+ = [0, \infty)$ , is called  $s$ -convex in the second sense if

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y)$$

for all  $x, y \in \mathbb{R}^+$ ,  $t \in [0, 1]$ ,  $s \in (0, 1]$  and these functions are denoted by  $f \in k_s^2$ . After that, Dragomir and Fitzpatrick [22] used this newly class of functions and proved the following H-H inequality:

$$2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{b-a} \int_a^b f(x) dx \leq \frac{f(a)+f(b)}{s+1}. \quad (1.2)$$

For more recent integral inequalities related to the class of  $s$ -convex functions and its generalizations via different integral operators, one can consult [11, 19, 20, 24, 25, 34, 36, 40].

On the other hand, several studies have been carried out in the domain of  $q$ -analysis, beginning with Euler, in order to achieve proficiency in mathematics that constructs quantum computing  $q$ -calculus, which is considered a relationship between physics and mathematics. It has a wide range of applications in mathematics, including combinatorics, simple hypergeometric functions, number theory, orthogonal polynomials, and other sciences, as well as mechanics, relativity theory, and quantum theory [27, 32]. Euler is thought to be the inventor of this significant branch of mathematics. He used the  $q$ -parameter in Newton's work on infinite series. Later, Jackson presented the  $q$ -calculus, which knew no limits calculus, in a methodical manner [26, 30]. In 1966, Al-Salam [10] introduced a  $q$ -analogue of the  $q$ -fractional integral and  $q$ -Riemann-Liouville fractional. Since then, the related research has gradually increased. In particular, in 2013, Tariboon and Ntouyas introduced  ${}_a D_q$ -difference operator and  $q_a$ -integral in [46]. In 2020, Bermudo et al. introduced the notion of  ${}^b D_q$  derivative and  $q^b$ -integral in [12].

Many integral inequalities have been studied using quantum integrals for various types of functions. For example, in [3, 6, 8, 9, 12–14, 31, 37, 43–45], the authors used  ${}_a D_q$ ,  ${}^b D_q$ -derivatives and  $q_a, q^b$ -integrals to prove H-H integral inequalities and their left-right estimates for convex and coordinated convex functions. In [38], Noor et al. presented a generalized version of quantum H-H integral inequalities. For generalized quasi-convex functions, Nwaeze et al. proved certain parameterized quantum integral inequalities in [39]. Khan et al. proved quantum H-H inequality using the green function in [35]. Budak et al. [15], Ali et al. [2, 4], Chu et al. [21] and Vivas-Cortez et al. [47] developed new quantum Simpson's and quantum Newton's type inequalities for convex and coordinated convex functions. For quantum Ostrowski's inequalities for convex and co-ordinated convex functions, one can consult [5, 7, 16].

Inspired by these ongoing studies, we offer a variant of quantum H-H inequality (1.2) and Ostrowski type inequalities for  $s$ -convex functions in the second sense. Since the  $s$ -convexity is the generalization of convexity, therefore the inequalities proved in this paper using the  $s$ -convexity are the generalization of already proved inequalities for convexity that is the main motivation of this paper.

The following is the structure of this paper: A brief overview of the concepts of  $q$ -calculus, as well as some related works, is given in Section 2. In Section 3, we show the relationship between the results presented here and comparable results in the literature by proving quantum H-H inequalities for  $s$ -convex functions in the second sense. Quantum Ostrowski type inequalities for  $s$ -convex functions in the second sense are presented in Section 4. Some applications to special means are given in Section 5. Section 6 concludes with some recommendations for future studies.

## 2. Preliminaries of $q$ -calculus and some inequalities

In this section, we recollect some formerly regarded concepts. Also, here and further we use  $q \in (0, 1)$  and the following notation (see [32]):

$$[n]_q = \frac{1 - q^n}{1 - q} = 1 + q + q^2 + \dots + q^{n-1}, \quad q \in (0, 1).$$

In [30], Jackson gave the  $q$ -Jackson integral from 0 to  $b$  as follows:

$$\int_0^b f(x) \, d_q x = (1 - q)b \sum_{n=0}^{\infty} q^n f(bq^n) \quad (2.1)$$

provided the sum converges absolutely.

**Definition 2.1.** [46] The  $q_a$ -derivative of a mapping  $f : [a, b] \rightarrow \mathbb{R}$  at  $x \in [a, b]$  is defined as:

$${}_a D_q f(x) = \frac{f(x) - f(qx + (1 - q)a)}{(1 - q)(x - a)}, \quad x \neq a. \quad (2.2)$$

If  $x = a$ , we define  ${}_a D_q f(a) = \lim_{x \rightarrow a} {}_a D_q f(x)$  if it exists and it is finite.

**Definition 2.2.** [12] The  $q^b$ -derivative of a mapping  $f : [a, b] \rightarrow \mathbb{R}$  at  $x \in [a, b]$  is defined as:

$${}^b D_q f(x) = \frac{f(qx + (1 - q)b) - f(x)}{(1 - q)(b - x)}, \quad x \neq b.$$

If  $x = b$ , we define  ${}^b D_q f(b) = \lim_{x \rightarrow b} {}^b D_q f(x)$  if it exists and it is finite.

**Definition 2.3.** [46] The  $q_a$ -integral of a mapping  $f : [a, b] \rightarrow \mathbb{R}$  is defined as:

$$\int_a^x f(t) \, {}_a d_q t = (1 - q)(x - a) \sum_{n=0}^{\infty} q^n f(q^n x + (1 - q^n)a),$$

where  $x \in [a, b]$ .

**Definition 2.4.** [12] The  $q^b$ -integral of a mapping  $f : [a, b] \rightarrow \mathbb{R}$  is defined as:

$$\int_x^b f(t) {}^b d_q t = (1-q)(b-x) \sum_{n=0}^{\infty} q^n f(q^n x + (1-q^n)b),$$

where  $x \in [a, b]$ .

In [12], Bermudo et al. established the following quantum H-H type inequality:

**Theorem 2.1.** For the convex mapping  $f : [a, b] \rightarrow \mathbb{R}$ , the following inequality holds

$$f\left(\frac{a+b}{2}\right) \leq \frac{1}{2(b-a)} \left[ \int_a^b f(x) {}_a d_q x + \int_a^b f(x) {}^b d_q x \right] \leq \frac{f(a) + f(b)}{2}. \quad (2.3)$$

In [16], Budak et al. proved the following Ostrowski inequality by using the concepts of quantum derivatives and integrals:

**Theorem 2.2.** Let  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function and  ${}^b D_q f$  and  ${}_a D_q f$  be two continuous and integrable functions on  $[a, b]$ . If  $|{}^b D_q f(t)|, |{}_a D_q f(t)| \leq M$  for all  $t \in [a, b]$ , then we have the following quantum Ostrowski inequality

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \leq \frac{qM}{(b-a)} \left[ \frac{(x-a)^2 + (b-x)^2}{[2]_q} \right] \quad (2.4)$$

### 3. Hermite-Hadamard inequalities

In this section, we prove H-H inequalities for  $s$ -convex functions in the second sense using the quantum integrals.

**Theorem 3.1.** Assume that the mapping  $f : \mathbb{R}^+ \rightarrow \mathbb{R}$  is  $s$ -convex in the second sense and  $a, b \in \mathbb{R}^+$  with  $a < b$ . If  $f \in L_1([a, b])$ , then the following inequality holds for  $s \in (0, 1]$ :

$$\begin{aligned} 2^{s-1} f\left(\frac{a+b}{2}\right) &\leq \frac{1}{2(b-a)} \left[ \int_a^b f(x) {}_a d_q x + \int_a^b f(x) {}^b d_q x \right] \\ &\leq \frac{f(a) + f(b)}{[s+1]_q}. \end{aligned} \quad (3.1)$$

*Proof.* As  $f$  is  $s$ -convex in the second sense on  $\mathbb{R}^+$  we have

$$f(tx + (1-t)y) \leq t^s f(x) + (1-t)^s f(y),$$

for all  $x, y \in \mathbb{R}^+$  and  $t \in [0, 1]$ .

Obverse that

$$2^s f\left(\frac{x+y}{2}\right) \leq f(x) + f(y). \quad (3.2)$$

We get the following, by putting  $x = tb + (1-t)a$  and  $y = ta + (1-t)b$  in (3.2)

$$2^s f\left(\frac{a+b}{2}\right) \leq f(tb + (1-t)a) + f(ta + (1-t)b).$$

From Definitions 2.3 and 2.4, we have

$$2^{s-1} f\left(\frac{a+b}{2}\right) \leq \frac{1}{2(b-a)} \left[ \int_a^b f(x) {}_a d_q x + \int_a^b f(x) {}^b d_q x \right]$$

and the first inequality in (3.1) is proved.

To proved the second inequality, we use the  $s$ -convexity and we have

$$f(tb + (1-t)a) \leq t^s f(b) + (1-t)^s f(a) \quad (3.3)$$

and

$$f(ta + (1-t)b) \leq t^s f(a) + (1-t)^s f(b). \quad (3.4)$$

By adding (3.3) and (3.4), from Definition 2.3 and 2.4, we have

$$\frac{1}{2(b-a)} \left[ \int_a^b f(x) {}_a d_q x + \int_a^b f(x) {}^b d_q x \right] \leq \frac{f(a) + f(b)}{[s+1]_q}$$

and the proof is completed.  $\square$

**Remark 3.1.** If we set  $s = 1$  in Theorem 3.1, then we recapture the inequality (2.3).

**Remark 3.2.** In Theorem 3.1, if we take the limit as  $q \rightarrow 1^-$ , then inequality (3.1) becomes the inequality (1.2).

#### 4. Ostrowski's inequalities

In this section, we prove Ostrowski's type inequalities for  $s$ -convex functions in the second sense. We use the following lemma to prove the new results.

**Lemma 4.1.** [16] Let  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$  be a function. If  ${}^b D_q f$  and  ${}_a D_q f$  are two continuous and integrable functions on  $[a, b]$ , then for all  $x \in [a, b]$  we have

$$\begin{aligned} & f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \\ &= \frac{q(x-a)^2}{b-a} \int_0^1 t {}_a D_q f(tx + (1-t)a) d_q t \\ &\quad - \frac{q(b-x)^2}{b-a} \int_0^1 t {}^b D_q f(tx + (1-t)b) d_q t. \end{aligned} \quad (4.1)$$

**Theorem 4.1.** Assume that the mapping  $f : I \subset \mathbb{R}^+ \rightarrow \mathbb{R}$  is differentiable and  $a, b \in I$  with  $a < b$ . If  $|{}_aD_q f|$  and  $|{}^bD_q f|$  are  $s$ -convex mappings in the second sense, then the following inequality holds:

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q(x-a)^2}{b-a} \left[ \frac{1}{[s+2]_q} |{}_aD_q f(x)| + \Theta_1 |{}_aD_q f(a)| \right] \\ & \quad + \frac{q(b-x)^2}{b-a} \left[ \frac{1}{[s+2]_q} |{}^bD_q f(x)| + \Theta_1 |{}^bD_q f(b)| \right] \end{aligned} \quad (4.2)$$

where

$$\Theta_1 = \int_0^1 t(1-t)^s d_q t.$$

*Proof.* From Lemma 4.1 and properties of the modulus, we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q(x-a)^2}{b-a} \int_0^1 t |{}_aD_q f(tx + (1-t)a)| d_q t + \frac{q(b-x)^2}{b-a} \int_0^1 t |{}^bD_q f(tx + (1-t)b)| d_q t. \end{aligned} \quad (4.3)$$

Since the mappings  $|{}_aD_q f|$  and  $|{}^bD_q f|$  are  $s$ -convex in the second sense, therefore

$$\begin{aligned} \int_0^1 t |{}_aD_q f(tx + (1-t)a)| d_q t & \leq \int_0^1 t^{s+1} |{}_aD_q f(x)| d_q t + \int_0^1 t(1-t)^s |{}_aD_q f(a)| d_q t \\ & = \frac{1}{[s+2]_q} |{}_aD_q f(x)| + \Theta_1 |{}_aD_q f(a)| \end{aligned} \quad (4.4)$$

and

$$\begin{aligned} \int_0^1 t |{}^bD_q f(tx + (1-t)b)| d_q t & \leq \int_0^1 t^{s+1} |{}^bD_q f(x)| d_q t + \int_0^1 t(1-t)^s |{}^bD_q f(b)| d_q t \\ & = \frac{1}{[s+2]_q} |{}^bD_q f(x)| + \Theta_1 |{}^bD_q f(b)|. \end{aligned} \quad (4.5)$$

We obtain the resultant inequality (4.2) by putting (4.4) and (4.5) in (4.3).  $\square$

**Remark 4.1.** If we set  $s = 1$  in Theorem 4.1, then we obtain the following inequality

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right|$$

$$\leq \frac{q}{(b-a)(1+q)(1+q+q^2)} \left[ (x-a)^2 \left( (1+q) \left| {}_a D_q f(x) \right| + q^2 \left| {}_a D_q f(a) \right| \right) \right. \\ \left. + (b-x)^2 \left( (1+q) \left| {}^b D_q f(x) \right| + q^2 \left| {}^b D_q f(b) \right| \right) \right]$$

which is given by Budak et al. in [16].

**Corollary 4.1.** If we assume  $|{}_a D_q f(x)|, |{}^b D_q f(x)| \leq M$  in Theorem 4.1, then we have following quantum Ostrowski's type inequality for  $s$ -convex functions in the second sense:

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ \leq \frac{Mq}{b-a} \left( \frac{1}{[s+2]_q} + \Theta_1 \right) \left[ (x-a)^2 + (b-x)^2 \right]. \quad (4.6)$$

**Remark 4.2.** If we set  $s = 1$  in Corollary 4.1, then we recapture inequality (2.4).

**Remark 4.3.** In Corollary 4.1, if we take the limit as  $q \rightarrow 1^-$ , then Corollary 4.1 reduces to [1, Theorem 2].

**Theorem 4.2.** Assume that the mapping  $f : I \subset \mathbb{R}^+ \rightarrow \mathbb{R}$  is differentiable and  $a, b \in I$  with  $a < b$ . If  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$ ,  $p_1 \geq 1$  are  $s$ -convex mappings in the second sense, then the following inequality holds:

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ \leq \frac{q}{b-a} \left( \frac{1}{[2]_q} \right)^{1-\frac{1}{p_1}} \left[ (x-a)^2 \left( \frac{1}{[s+2]_q} \left| {}_a D_q f(x) \right|^{p_1} + \Theta_1 \left| {}_a D_q f(a) \right|^{p_1} \right)^{\frac{1}{p_1}} \right. \\ \left. + (b-x)^2 \left( \frac{1}{[s+2]_q} \left| {}^b D_q f(x) \right|^{p_1} + \Theta_1 \left| {}^b D_q f(b) \right|^{p_1} \right)^{\frac{1}{p_1}} \right]. \quad (4.7)$$

*Proof.* From Lemma 4.1, using properties of the modulus and power mean inequality, we have

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ \leq \frac{q(x-a)^2}{b-a} \int_0^1 t \left| {}_a D_q f(tx + (1-t)a) \right| d_q t + \frac{q(b-x)^2}{b-a} \int_0^1 t \left| {}^b D_q f(tx + (1-t)b) \right| d_q t \\ \leq \frac{q(x-a)^2}{b-a} \left( \int_0^1 t d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t \left| {}_a D_q f(tx + (1-t)a) \right|^{p_1} d_q t \right)^{\frac{1}{p_1}}$$

$$+ \frac{q(b-x)^2}{b-a} \left( \int_0^1 t d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t | {}^b D_q f(tx + (1-t)b) |^{p_1} d_q t \right)^{\frac{1}{p_1}}. \quad (4.8)$$

Since the mappings  $| {}_a D_q f |^{p_1}$  and  $| {}^b D_q f |^{p_1}$  are  $s$ -convex in the second sense, therefore

$$\begin{aligned} & \left( \int_0^1 t d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t | {}_a D_q f(tx + (1-t)a) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[2]_q} \right)^{1-\frac{1}{p_1}} \left( \frac{1}{[s+2]_q} | {}_a D_q f(x) |^{p_1} + \Theta_1 | {}_a D_q f(a) |^{p_1} \right)^{\frac{1}{p_1}} \end{aligned} \quad (4.9)$$

and

$$\begin{aligned} & \left( \int_0^1 t d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t | {}^b D_q f(tx + (1-t)b) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[2]_q} \right)^{1-\frac{1}{p_1}} \left( \frac{1}{[s+2]_q} | {}^b D_q f(x) |^{p_1} + \Theta_1 | {}^b D_q f(b) |^{p_1} \right)^{\frac{1}{p_1}}. \end{aligned} \quad (4.10)$$

We obtain the resultant inequality (4.7) by putting (4.9) and (4.10) in (4.8).  $\square$

**Remark 4.4.** If we set  $s = 1$  in Theorem 4.2, then we obtain the following inequality

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q}{(b-a)[2]_q} \left[ (x-a)^2 \left( \frac{[2]_q | {}_a D_q f(x) |^{p_1} + q^2 | {}_a D_q f(a) |^{p_1}}{[3]_q} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + (b-x)^2 \left( \frac{[2]_q | {}^b D_q f(x) |^{p_1} + q^2 | {}^b D_q f(b) |^{p_1}}{[3]_q} \right)^{\frac{1}{p_1}} \right] \end{aligned}$$

which is proved by Budak et al. in [16].

**Corollary 4.2.** If we assume  $| {}_a D_q f(x), | {}_a D_q f(a) | \leq M$  in Theorem 4.2, then we have following quantum Ostrowski's type inequality for  $s$ -convex functions in the second sense:

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{Mq}{b-a} \left( \frac{1}{[2]_q} \right)^{1-\frac{1}{p_1}} \left( \frac{1}{[s+2]_q} + \Theta_1 \right)^{\frac{1}{p_1}} \left[ (x-a)^2 + (b-x)^2 \right]. \end{aligned}$$



**Remark 4.5.** In Corollary 4.2, if we take the limit as  $q \rightarrow 1^-$ , then Corollary 4.2 reduces to [1, Theorem 4].

**Theorem 4.3.** Assume that the mapping  $f : I \subset \mathbb{R}^+ \rightarrow \mathbb{R}$  is differentiable and  $a, b \in I$  with  $a < b$ . If  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$ ,  $p_1 > 1$  are  $s$ -convex mappings in the second sense, then the following inequality holds:

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q}{b-a} \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left[ (x-a)^2 \left( \frac{1}{[s+1]_q} \left( |{}_a D_q f(x)|^{p_1} + |{}_a D_q f(a)|^{p_1} \right) \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + (b-x)^2 \left( \frac{1}{[s+1]_q} \left( |{}^b D_q f(x)|^{p_1} + |{}^b D_q f(b)|^{p_1} \right) \right)^{\frac{1}{p_1}} \right], \end{aligned} \quad (4.11)$$

where  $\frac{1}{r_1} + \frac{1}{p_1} = 1$ .

*Proof.* From Lemma 4.1, using properties of the modulus and Hölder's inequality, we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q(x-a)^2}{b-a} \int_0^1 t |{}_a D_q f(tx + (1-t)a)| d_q t + \frac{q(b-x)^2}{b-a} \int_0^1 t |{}^b D_q f(tx + (1-t)b)| d_q t \\ & \leq \frac{q(x-a)^2}{b-a} \left( \int_0^1 t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 |{}_a D_q f(tx + (1-t)a)|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \quad + \frac{q(b-x)^2}{b-a} \left( \int_0^1 t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 |{}^b D_q f(tx + (1-t)b)|^{p_1} d_q t \right)^{\frac{1}{p_1}}. \end{aligned} \quad (4.12)$$

Since the mappings  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$  are  $s$ -convex in the second sense, therefore

$$\begin{aligned} & \left( \int_0^1 t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 |{}_a D_q f(tx + (1-t)a)|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left( \frac{1}{[s+1]_q} \left( |{}_a D_q f(x)|^{p_1} + |{}_a D_q f(a)|^{p_1} \right) \right)^{\frac{1}{p_1}} \end{aligned} \quad (4.13)$$

and

$$\left( \int_0^1 t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 |{}^b D_q f(tx + (1-t)b)|^{p_1} d_q t \right)^{\frac{1}{p_1}}$$

$$\leq \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left( \frac{1}{[s + 1]_q} \left( | {}^b D_q f(x) |^{p_1} + | {}^b D_q f(b) |^{p_1} \right) \right)^{\frac{1}{p_1}}. \quad (4.14)$$

We obtain the resultant inequality (4.11) by putting (4.13) and (4.14) in (4.12).  $\square$

**Remark 4.6.** If we set  $s = 1$  in Theorem 4.3, then we obtain the following inequality

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q}{b-a} \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left[ (x-a)^2 \left( \frac{| {}_a D_q f(x) |^{p_1} + q | {}_a D_q f(a) |^{p_1}}{[2]_q} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + (b-x)^2 \left( \frac{| {}^b D_q f(x) |^{p_1} + q | {}^b D_q f(b) |^{p_1}}{[2]_q} \right)^{\frac{1}{p_1}} \right] \end{aligned}$$

which is proved by Budak et al. in [16].

**Corollary 4.3.** If we assume  $| {}_a D_q f(x) |, | {}^b D_q f(a) | \leq M$  in Theorem 4.3, then we have the following quantum Ostrowski's type inequality for  $s$ -convex functions in the second sense:

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{Mq}{b-a} \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left( \frac{2}{[s + 1]_q} \right)^{\frac{1}{p_1}} \left[ (x-a)^2 + (b-x)^2 \right]. \quad (4.15) \end{aligned}$$

**Remark 4.7.** In Corollary 4.3, if we set  $s = 1$ , then we recapture the following inequality

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{qM}{b-a} \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \left[ (x-a)^2 + (b-x)^2 \right] \end{aligned}$$

which is obtained by Budak et al. in [16].

**Remark 4.8.** In Corollary 4.3, if we take the limit as  $q \rightarrow 1^-$ , then Corollary 4.3 reduces to [1, Theorem 3].

**Theorem 4.4.** Assume that the mapping  $f : I \subset \mathbb{R}^+ \rightarrow \mathbb{R}$  is differentiable and  $a, b \in I$  with  $a < b$ . If  $| {}_a D_q f |^{p_1}$  and  $| {}^b D_q f |^{p_1}$ ,  $p_1 \geq 1$  are  $s$ -convex mappings in the second sense, then the following inequality holds:

$$\left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right|$$

$$\begin{aligned}
&\leq \frac{q}{(b-a)} \left( \frac{q^2}{[2]_q [3]_q} \right)^{1-\frac{1}{p_1}} \\
&\quad \times \left[ (x-a)^2 \left( \left( \frac{1}{[s+2]_q} - \frac{1}{[s+3]_q} \right) |{}_a D_q f(x)|^{p_1} + \Theta_2 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}} \right. \\
&\quad \left. + (b-x)^2 \left( \left( \frac{1}{[s+2]_q} - \frac{1}{[s+3]_q} \right) |{}^b D_q f(x)|^{p_1} + \Theta_2 |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \right] \\
&\quad + \frac{q}{(b-a)} \left( \frac{1}{[3]_q} \right)^{1-\frac{1}{p_1}} \left[ (x-a)^2 \left( \frac{1}{[s+3]_q} |{}_a D_q f(x)|^{p_1} + \Theta_3 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}} \right. \\
&\quad \left. + (b-x)^2 \left( \frac{1}{[s+3]_q} |{}^b D_q f(x)|^{p_1} + \Theta_3 |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \right], \tag{4.16}
\end{aligned}$$

where

$$\begin{aligned}
\Theta_2 &= \int_0^1 t(1-t)^{s+1} d_q t, \\
\Theta_3 &= \int_0^1 t^2(1-t)^s d_q t.
\end{aligned}$$

*Proof.* From Lemma 4.1, using properties of the modulus and improved power mean inequality (see [33]), we have

$$\begin{aligned}
&\left\| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right\| \\
&\leq \frac{q(x-a)^2}{b-a} \int_0^1 t |{}_a D_q f(tx + (1-t)a)| d_q t + \frac{q(b-x)^2}{b-a} \int_0^1 t |{}^b D_q f(tx + (1-t)b)| d_q t \\
&\leq \frac{q(x-a)^2}{b-a} \left( \int_0^1 t(1-t) d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t(1-t) |{}_a D_q f(tx + (1-t)a)|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\quad + \frac{q(x-a)^2}{b-a} \left( \int_0^1 t^2 d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t^2 |{}_a D_q f(tx + (1-t)a)|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\quad + \frac{q(b-x)^2}{b-a} \left( \int_0^1 t(1-t) d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t(1-t) |{}^b D_q f(tx + (1-t)b)|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\quad + \frac{q(b-x)^2}{b-a} \left( \int_0^1 t^2 d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t^2 |{}^b D_q f(tx + (1-t)b)|^{p_1} d_q t \right)^{\frac{1}{p_1}}. \tag{4.17}
\end{aligned}$$

Since the mappings  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$  are  $s$ -convex in the second sense, therefore

$$\begin{aligned} & \left( \int_0^1 t(1-t) d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t(1-t) |{}_a D_q f(tx + (1-t)a|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{q^2}{[2]_q [3]_q} \right)^{1-\frac{1}{p_1}} \left( \left( \frac{1}{[s+2]_q} - \frac{1}{[s+3]_q} \right) |{}_a D_q f(x)|^{p_1} + \Theta_2 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}}, \end{aligned} \quad (4.18)$$

$$\begin{aligned} & \left( \int_0^1 t^2 d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t^2 |{}_a D_q f(tx + (1-t)a|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[3]_q} \right)^{1-\frac{1}{p_1}} \left( \frac{1}{[s+3]_q} |{}_a D_q f(x)|^{p_1} + \Theta_3 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}}, \end{aligned} \quad (4.19)$$

$$\begin{aligned} & \left( \int_0^1 t(1-t) d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t(1-t) |{}^b D_q f(tx + (1-t)b|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{q^2}{[2]_q [3]_q} \right)^{1-\frac{1}{p_1}} \left( \left( \frac{1}{[s+2]_q} - \frac{1}{[s+3]_q} \right) |{}^b D_q f(x)|^{p_1} + \Theta_2 |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \end{aligned} \quad (4.20)$$

and

$$\begin{aligned} & \left( \int_0^1 t^2 d_q t \right)^{1-\frac{1}{p_1}} \left( \int_0^1 t^2 |{}^b D_q f(tx + (1-t)b|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[3]_q} \right)^{1-\frac{1}{p_1}} \left( \frac{1}{[s+3]_q} |{}^b D_q f(x)|^{p_1} + \Theta_3 |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}}. \end{aligned} \quad (4.21)$$

We obtain the resultant inequality (4.16) by putting (4.18), (4.19), (4.20) and (4.21) in (4.17).  $\square$

**Corollary 4.4.** *If we assume  $|{}_a D_q f(x)|, |{}^b D_q f(x)| \leq M$  in Theorem 4.4, then we have the following quantum Ostrowski's type inequality for  $s$ -convex functions in the second sense:*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{qM}{(b-a)} \left[ \left( \frac{q^2}{[2]_q [3]_q} \right)^{1-\frac{1}{p_1}} [(x-a)^2 + (b-x)^2] \left( \frac{1}{[s+2]_q} - \frac{1}{[s+3]_q} + \Theta_2 \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + \left( \frac{1}{[3]_q} \right)^{1-\frac{1}{p_1}} [(x-a)^2 + (b-x)^2] \left( \frac{1}{[s+3]_q} + \Theta_3 \right)^{\frac{1}{p_1}} \right]. \end{aligned} \quad (4.22)$$

**Corollary 4.5.** *In Corollary 4.4, if we set  $s = 1$ , then we obtain the following new Ostrowski type inequality:*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{qM}{(b-a)} \left[ \left( \frac{q^2}{[2]_q [3]_q} \right)^{1-\frac{1}{p_1}} [(x-a)^2 + (b-x)^2] \left( \frac{q^2}{[2]_q [3]_q} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + \left( \frac{1}{[3]_q} \right)^{1-\frac{1}{p_1}} [(x-a)^2 + (b-x)^2] \left( \frac{1}{[3]_q} \right)^{\frac{1}{p_1}} \right]. \end{aligned}$$

**Theorem 4.5.** *Assume that the mapping  $f : I \subset \mathbb{R}^+ \rightarrow \mathbb{R}$  is differentiable and  $a, b \in I$  with  $a < b$ . If  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$ ,  $p_1 > 1$  are  $s$ -convex mappings in the second sense, then the following inequality holds:*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q}{b-a} \left( \frac{1}{[r_1+1]_q} - \frac{1}{[r_1+2]_q} \right)^{\frac{1}{r_1}} \\ & \quad \times \left[ (x-a)^2 \left( \left( \frac{1}{[s+1]_q} - \frac{1}{[s+2]_q} \right) |{}_a D_q f(x)|^{p_1} + \frac{1}{[s+2]_q} |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + (b-x)^2 \left( \left( \frac{1}{[s+1]_q} - \frac{1}{[s+2]_q} \right) |{}^b D_q f(x)|^{p_1} + \frac{1}{[s+2]_q} |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \right] \\ & \quad + \frac{q}{b-a} \left( \frac{1}{[r_1+1]_q} \right)^{\frac{1}{r_1}} \\ & \quad \times \left[ (x-a)^2 \left( \frac{1}{[s+2]_q} |{}_a D_q f(x)|^{p_1} + \Theta_1 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + (b-x)^2 \left( \frac{1}{[s+2]_q} |{}^b D_q f(x)|^{p_1} + \Theta_1 |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \right], \end{aligned} \tag{4.23}$$

where  $\frac{1}{r_1} + \frac{1}{p_1} = 1$ .

*Proof.* From Lemma 4.1, using properties of the modulus and Hölder's İşcan inequality (see [29]), we have

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{q(x-a)^2}{b-a} \int_0^1 t |{}_a D_q f(tx + (1-t)a)| d_q t + \frac{q(b-x)^2}{b-a} \int_0^1 t |{}^b D_q f(tx + (1-t)b)| d_q t \end{aligned}$$

$$\begin{aligned}
&\leq \frac{q(x-a)^2}{b-a} \left( \int_0^1 (1-t)t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 (1-t) | {}_a D_q f(tx + (1-t)a) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&+ \frac{q(x-a)^2}{b-a} \left( \int_0^1 t^{r_1+1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 t | {}_a D_q f(tx + (1-t)a) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&+ \frac{q(b-x)^2}{b-a} \left( \int_0^1 (1-t)t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 (1-t) | {}^b D_q f(tx + (1-t)b) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&+ \frac{q(b-x)^2}{b-a} \left( \int_0^1 t^{r_1+1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 t | {}^b D_q f(tx + (1-t)b) |^{p_1} d_q t \right)^{\frac{1}{p_1}}. \tag{4.24}
\end{aligned}$$

Since the mappings  $|{}_a D_q f|^{p_1}$  and  $|{}^b D_q f|^{p_1}$  are  $s$ -convex in the second sense, therefore

$$\begin{aligned}
&\left( \int_0^1 (1-t)t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 (1-t) | {}_a D_q f(tx + (1-t)a) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\leq \left( \frac{1}{[r_1+1]_q} - \frac{1}{[r_1+2]_q} \right)^{\frac{1}{r_1}} \\
&\times \left( \left( \frac{1}{[s+1]_q} - \frac{1}{[s+2]_q} \right) |{}_a D_q f(x)|^{p_1} + \frac{1}{[s+2]_q} |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}}, \tag{4.25}
\end{aligned}$$

$$\begin{aligned}
&\left( \int_0^1 t^{r_1+1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 t | {}_a D_q f(tx + (1-t)a) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\leq \left( \frac{1}{[r_1+2]_q} \right)^{\frac{1}{r_1}} \left( \frac{1}{[s+2]_q} |{}_a D_q f(x)|^{p_1} + \Theta_1 |{}_a D_q f(a)|^{p_1} \right)^{\frac{1}{p_1}}, \tag{4.26}
\end{aligned}$$

$$\begin{aligned}
&\left( \int_0^1 (1-t)t^{r_1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 (1-t) | {}^b D_q f(tx + (1-t)b) |^{p_1} d_q t \right)^{\frac{1}{p_1}} \\
&\leq \left( \frac{1}{[r_1+1]_q} - \frac{1}{[r_1+2]_q} \right)^{\frac{1}{r_1}} \\
&\times \left( \left( \frac{1}{[s+1]_q} - \frac{1}{[s+2]_q} \right) |{}^b D_q f(x)|^{p_1} + \frac{1}{[s+2]_q} |{}^b D_q f(b)|^{p_1} \right)^{\frac{1}{p_1}} \tag{4.27}
\end{aligned}$$

and

$$\begin{aligned} & \left( \int_0^1 t^{r_1+1} d_q t \right)^{\frac{1}{r_1}} \left( \int_0^1 t \left| {}^b D_q f(tx + (1-t)b) \right|^{p_1} d_q t \right)^{\frac{1}{p_1}} \\ & \leq \left( \frac{1}{[r_1 + 2]_q} \right)^{\frac{1}{r_1}} \left( \frac{1}{[s + 2]_q} \left| {}^b D_q f(x) \right|^{p_1} + \Theta_1 \left| {}^b D_q f(b) \right|^{p_1} \right)^{\frac{1}{p_1}}. \end{aligned} \quad (4.28)$$

We obtain the resultant inequality (4.23) by putting (4.25), (4.26), (4.27) and (4.28) in (4.24).  $\square$

**Corollary 4.6.** *If we assume  $|{}_a D_q f(x)|, |{}^b D_q f(x)| \leq M$  in Theorem 4.5, then we have the following quantum Ostrowski's type inequality for  $s$ -convex functions in the second sense:*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{Mq}{b-a} \\ & \quad \times \left[ \left( \frac{1}{[r_1 + 1]_q} - \frac{1}{[r_1 + 2]_q} \right)^{\frac{1}{r_1}} [(x-a)^2 + (b-x)^2] \left( \frac{1}{[s + 1]_q} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} [(x-a)^2 + (b-x)^2] \left( \frac{1}{[s + 2]_q} + \Theta_1 \right)^{\frac{1}{p_1}} \right]. \end{aligned} \quad (4.29)$$

**Corollary 4.7.** *In Corollary 4.6, if we use  $s = 1$ , then we obtain the following new Ostrowski type inequality:*

$$\begin{aligned} & \left| f(x) - \frac{1}{b-a} \left[ \int_a^x f(t) {}_a d_q t + \int_x^b f(t) {}^b d_q t \right] \right| \\ & \leq \frac{Mq}{b-a} \left( \frac{1}{[2]_q} \right)^{\frac{1}{p_1}} [(x-a)^2 + (b-x)^2] \\ & \quad \times \left[ \left( \frac{1}{[r_1 + 1]_q} - \frac{1}{[r_1 + 2]_q} \right)^{\frac{1}{r_1}} + \left( \frac{1}{[r_1 + 1]_q} \right)^{\frac{1}{r_1}} \right]. \end{aligned}$$

## 5. Applications to special means

For arbitrary positive numbers  $\kappa_1, \kappa_2$  ( $\kappa_1 \neq \kappa_2$ ), we consider the means as follows:

1. The arithmetic mean

$$\mathcal{A} = \mathcal{A}(\kappa_1, \kappa_2) = \frac{\kappa_1 + \kappa_2}{2}.$$

2. The logarithmic mean

$$\mathcal{L}_p^p = \mathcal{L}_p^p(\kappa_1, \kappa_2) = \frac{\kappa_2^{p+1} - \kappa_1^{p+1}}{(p+1)(\kappa_2 - \kappa_1)}.$$

**Proposition 5.1.** For  $0 < a < b$  and  $0 < q < 1$ , the following inequality is true:

$$\begin{aligned} & \left| \frac{1}{s+1} [\mathcal{A}^{s+1}(a, b) - \mathcal{A}(\mathbb{k}_1, \mathbb{k}_2)] \right| \\ & \leq \frac{q(b-a)}{2} \left[ \frac{1}{[s+2]_q} \left\{ \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)a, \frac{a+b}{2} \right) \right. \right. \\ & \quad \left. \left. + \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)b, \frac{a+b}{2} \right) \right\} + 2\Theta_1 \mathcal{A}(a^s, b^s) \right], \end{aligned}$$

where

$$\begin{aligned} \mathbb{k}_1 &= (1-q) \sum_{n=0}^{\infty} q^n \left( q^n \frac{a+b}{2} + (1-q^n)a \right)^{s+1}, \\ \mathbb{k}_2 &= (1-q) \sum_{n=0}^{\infty} q^n \left( q^n \frac{a+b}{2} + (1-q^n)b \right)^{s+1}. \end{aligned}$$

*Proof.* The inequality (4.2) in Theorem 4.1 with  $x = \frac{a+b}{2}$  for  $f(x) = \frac{x^{s+1}}{s+1}$ , where  $x > 0$  and  $s \in (0, 1)$  leads to this conclusion.  $\square$

**Proposition 5.2.** For  $0 < a < b$  and  $0 < q < 1$ , the following inequality is true:

$$\begin{aligned} & \left| \frac{1}{s+1} [\mathcal{A}^{s+1}(a, b) - \mathcal{A}(\mathbb{k}_1, \mathbb{k}_2)] \right| \\ & \leq \frac{Mq(b-a)}{2} \left[ \frac{1}{[s+2]_q} + \Theta_1 \right]. \end{aligned}$$

*Proof.* The inequality (4.6) in Corollary 4.1 with  $x = \frac{a+b}{2}$  for  $f(x) = \frac{x^{s+1}}{s+1}$ , where  $x > 0$  and  $s \in (0, 1)$  leads to this conclusion.  $\square$

**Proposition 5.3.** For  $0 < a < b$  and  $0 < q < 1$ , the following inequality is true:

$$\begin{aligned} & \left| \frac{1}{s+1} [\mathcal{A}^{s+1}(a, b) - \mathcal{A}(\mathbb{k}_1, \mathbb{k}_2)] \right| \\ & \leq \frac{q(b-a)}{2} \left( \frac{1}{[2]_q} \right)^{1-\frac{1}{p_1}} \left[ \left( \frac{1}{[s+2]_q} \left| \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)a, \frac{a+b}{2} \right) \right|^{p_1} + \Theta_1 |a^s|^{p_1} \right)^{\frac{1}{p_1}} \right. \\ & \quad \left. + \left( \frac{1}{[s+2]_q} \left| \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)b, \frac{a+b}{2} \right) \right|^{p_1} + \Theta_1 |b^s|^{p_1} \right)^{\frac{1}{p_1}} \right]. \end{aligned}$$

*Proof.* The inequality (4.7) in Theorem 4.2 with  $x = \frac{a+b}{2}$  for  $f(x) = \frac{x^{s+1}}{s+1}$ , where  $x > 0$  and  $s \in (0, 1)$  leads to this conclusion.  $\square$

**Proposition 5.4.** For  $0 < a < b$  and  $0 < q < 1$ , the following inequality is true:

$$\left| \frac{1}{s+1} [\mathcal{A}^{s+1}(a, b) - \mathcal{A}(\mathbb{k}_1, \mathbb{k}_2)] \right|$$



$$\leq \frac{q(b-a)}{2} \left( \frac{1}{[r_1+1]_q} \right)^{\frac{1}{r_1}} \left[ \left( \frac{1}{[s+1]_q} \left( \left| \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)a, \frac{a+b}{2} \right) \right|^{p_1} + |a^s|^{p_1} \right) \right)^{\frac{1}{p_1}} \right. \\ \left. + \left( \frac{1}{[s+1]_q} \left( \left| \mathcal{L}_s^s \left( q \frac{a+b}{2} + (1-q)b, \frac{a+b}{2} \right) \right|^{p_1} + |b^s|^{p_1} \right) \right)^{\frac{1}{p_1}} \right].$$

*Proof.* The inequality (4.11) in Theorem 4.3 with  $x = \frac{a+b}{2}$  for  $f(x) = \frac{x^{s+1}}{s+1}$ , where  $x > 0$  and  $s \in (0, 1)$  leads to this conclusion.  $\square$

**Proposition 5.5.** For  $0 < a < b$  and  $0 < q < 1$ , the following inequality is true:

$$\left| \frac{1}{s+1} [\mathcal{A}^{s+1}(a, b) - \mathcal{A}(\mathbb{k}_1, \mathbb{k}_2)] \right| \\ \leq \frac{Mq(b-a)}{2} \left( \frac{1}{[r_1+1]_q} \right)^{\frac{1}{r_1}} \left( \frac{2}{[s+1]_q} \right)^{\frac{1}{p_1}}.$$

*Proof.* The inequality (4.15) in Corollary 4.3 with  $x = \frac{a+b}{2}$  for  $f(x) = \frac{x^{s+1}}{s+1}$ , where  $x > 0$  and  $s \in (0, 1)$  leads to this conclusion.  $\square$

## 6. Conclusions

In this investigation, Hermite-Hadamard and Ostrowski type inequalities for  $s$ -convex mappings in the second sense are derived, by applying quantum integrals. It is also showed that the results established in this paper are potential generalization of the existing comparable results in the literature. As future directions, one can find similar inequalities for co-ordinated  $s$ -convex functions in the second sense.

## Acknowledgments

This research was funded by King Mongkut's University of Technology North Bangkok. Contract no. KMUTNB-61-KNOW-015. The work of second author is partially supported by the National Natural Science Foundation of China (Grant No. 11971241).

## Conflict of interest

The authors declare that they have no competing interest.

## References

1. M. Alomari, M. Darus, S. S. Dragomir, Ostrowski type inequalities for functions whose derivatives are  $s$ -convex in the second sense, *Appl. Math. Lett.*, **23** (2010), 1071–1076.
2. M. A. Ali, H. Budak, Z. Zhang, H. Yildirim, Some new Simpson's type inequalities for co-ordinated convex functions in quantum calculus, *Math. Meth. Appl. Sci.*, **44** (2021), 4515–4540.

3. M. A. Ali, H. Budak, M. Abbas, Y. M. Chu, Quantum Hermite–Hadamard-type inequalities for functions with convex absolute values of second  $q^b$ -derivatives, *Adv. Differ. Equ.*, **2021** (2021), 7.
4. M. A. Ali, M. Abbas, H. Budak, P. Agarwal, G. Murtaza, Y. M. Chu, New quantum boundaries for quantum Simpson’s and quantum Newton’s type inequalities for preinvex functions, *Adv. Differ. Equ.*, **2021** (2021), 64.
5. M. A. Ali, Y. M. Chu, H. Budak, A. Akkurt, H. Yildirim, Quantum variant of Montgomery identity and Ostrowski-type inequalities for the mappings of two variables, *Adv. Differ. Equ.*, **2021** (2021), 25.
6. M. A. Ali, N. Alp, H. Budak, Y. M. Chu, Z. Zhang, On some new quantum midpoint type inequalities for twice quantum differentiable convex functions, *Open Math.*, 2021, in press.
7. M. A. Ali, H. Budak, A. Akkurt, Y. M. Chu, Quantum Ostrowski type inequalities for twice quantum differentiable functions in quantum calculus, *Open Math.*, 2021, in press.
8. N. Alp, M. Z. Sarikaya, M. Kunt, İ. İşcan,  $q$ -Hermite Hadamard inequalities and quantum estimates for midpoint type inequalities via convex and quasi-convex functions, *J. King Saud Univ. Sci.*, **30** (2018), 193–203.
9. N. Alp, M. Z. Sarikaya, Hermite Hadamard’s type inequalities for co-ordinated convex functions on quantum integral, *Appl. Math. E-Notes*, **20** (2020), 341–356.
10. W. Al-Salam, Some fractional  $q$ -integrals and  $q$ -derivatives, *Proc. Edinburgh Math. Soc.*, **15** (1966), 135–140.
11. B. Bayraktar, V. C. Kudaev, Some new integral inequalities for  $(s, m)$ -convex and  $(\alpha, m)$ -convex functions, *Bull. Karaganda Univ. Math.*, **94** (2019), 15–25.
12. S. Bermudo, P. Kórus, J. N. Valdés, On  $q$ -Hermite-Hadamard inequalities for general convex functions, *Acta Math. Hung.*, **162** (2020), 364–374.
13. H. Budak, Some trapezoid and midpoint type inequalities for newly defined quantum integrals, *Proyecciones*, **40** (2021), 199–215.
14. H. Budak, M. A. Ali, M. Tarhanaci, Some new quantum Hermite-Hadamard-like inequalities for coordinated convex functions, *J. Optim. Theory Appl.*, **186** (2020), 899–910.
15. H. Budak, S. Erden, M. A. Ali, Simpson and Newton type inequalities for convex functions via newly defined quantum integrals, *Math. Meth. Appl. Sci.*, **44** (2020), 378–390.
16. H. Budak, M. A. Ali, N. Alp, Y. M. Chu, Quantum Ostrowski type integral inequalities, *J. Math. Inequal.*, 2021, in press.
17. S. I. Butt, S. Yousuf, A. O. Akdemir, M. A. Dokuyucu, New Hadamard-type integral inequalities via a general form of fractional integral operators, *Chaos Solition Fract.*, **148** (2021), 111025.
18. S. I. Butt, A. O. Akdemir, N. Nadeem, N. Malaiki, İ. İşcan, T. Abdeljawad,  $(m, n)$ -Harmonically polynomial convex functions and some Hadamard type inequalities on the coordinates, *AIMS Math.*, **6** (2021), 4677–4690.
19. M. Çakmak, Refinements of Bullen-type inequalities for  $s$ -convex functions via Riemann–Liouville fractional integrals involving Gauss hypergeometric function, *J. Interdiscip. Math.*, **22** (2019), 975–989.

20. Y. M. Chu, S. Rashid, T. Abdeljawad, A. Khalid, H. Kalsoom, On new generalized unified bounds via generalized exponentially harmonically  $s$ -convex functions on fractal sets. *Adv. Differ. Equ.*, **2021** (2021), 1–33.
21. Y. M. Chu, A. Rauf, S. Rashid, S. Batool, Y. S. Hamed, Quantum estimates in two variable forms for Simpson-type inequalities considering generalized  $\Psi$ -convex functions with applications, *Open Phy.*, **19** (2021), 305–326.
22. S. S. Dragomir, S. Fitzpatrick, The Hadamard inequalities for  $s$ -convex functions in the second sense, *Demonstr. Math.*, **32** (1999), 68–696.
23. S. S. Dragomir, C. E. M. Pearce, *Selected topics on Hermite-Hadamard inequalities and applications*, RGMIA Monographs, Victoria University, 2000.
24. T. S. Du, Y. J. Li, Z. Q. Yang, A generalization of Simpson's inequality via differentiable mapping using extended  $(s, m)$ -convex functions, *Appl. Math. Comput.*, **293** (2017), 358–369.
25. T. S. Du, Y. Luo, B. Yu, Certain quantum estimates on the parameterized integral inequalities and their applications, *J. Math. Inequal.*, **15** (2021), 201–228.
26. T. Ernst, *The history of  $q$ -calculus and new method*, Sweden: Department of Mathematics, Uppsala University, 2000.
27. T. Ernst, *A comprehensive treatment of  $q$ -calculus*, Springer, Basel, 2012.
28. H. Hudzik, L. Maligranda, Some remarks on  $s$ -convex functions, *Aequationes Math.*, **48** (1994), 100–111.
29. İ. İşcan, New refinements for integral and sum forms of Hölder inequality, *J. Inequal. Appl.*, **2019** (2019), 1–11.
30. F. H. Jackson, On a  $q$ -definite integrals. *Quarterly J. Pure Appl. Math.*, **41** (1910), 193–203.
31. S. Jhathanam, T. Jessada, S. K. Ntouyas, K. Nonlaopon, On  $q$ -Hermite-Hadamard inequalities for differentiable convex functions, *Mathematics*, **7** (2019), 632.
32. V. Kac, P. Cheung, *Quantum calculus*, Springer, New York, 2002.
33. M. Kadakal, İ. İşcan, H. Kadakal, K. Bekar, On improvements of some integral inequalities, *Researchgate*, Preprint.
34. M. A. Khan, Y. M. Chu, T. U. Khan, J. Khan, Some new inequalities of Hermite–Hadamard type for  $s$ -convex functions with applications, *Open Math.*, **15** (2017), 1414–1430.
35. M. A. Khan, M. Noor, E. R. Nwaeze, Y. M. Chu, Quantum Hermite–Hadamard inequality by means of a Green function, *Adv. Differ. Equ.*, **2020** (2020), 99.
36. P. Kórus, An extension of the Hermite–Hadamard inequality for convex and  $s$ -convex functions, *Aequat. Math.*, **93** (2019), 527–534.
37. M. A. Noor, K. I. Noor, M. U. Awan, Some quantum estimates for Hermite-Hadamard inequalities, *Appl. Math. Comput.*, **251** (2015), 675–679.
38. M. A. Noor, K. I. Noor, M. U. Awan, Some quantum integral inequalities via preinvex functions, *Appl. Math. Comput.*, **269** (2015), 242–251.
39. E. R. Nwaeze, A. M. Tameru, New parameterized quantum integral inequalities via  $\eta$ -quasiconvexity, *Adv. Differ. Equ.*, **2019** (2019), 425.

40. M. E. Özdemir, S. I. Butt, B. Bayraktar, J. Nasir, Several integral inequalities for  $(\alpha, s, m)$ -convex functions, *AIMS Math.*, **5** (2020), 3906–3921.
41. J. E. Pečarić, F. Proschan, Y. L. Tong, *Convex functions, partial orderings and statistical applications*, Academic Press, Boston, 1992.
42. S. Rashid, A. Khalid, O. Bazighifan, G. I. Oros, New modifications of integral inequalities via  $P$ -convexity pertaining to fractional calculus and their applications, *Mathematics*, **9** (2021), 1753.
43. S. Rashid, S. Parveen, H. Ahmad, Y. M. Chu, New quantum integral inequalities for some new classes of generalized  $\psi$ -convex functions and their scope in physical systems, *Open Phy.*, **19** (2021), 35–50.
44. S. Rashid, Z. Hammouch, R. Ashraf, D. Baleanu, K. S. Nasir, New quantum estimates in the setting of fractional calculus theory, *Adv. Differ. Equ.*, **2020** (2020), 1–17.
45. S. Rashid, S. I. Butt, S. Kanwal, H. Ahmed, M. K. Wang, Quantum integral inequalities with respect to Raina's function via coordinated generalized  $\eta$ -convex functions with applications, *J. Funct. Spaces*, **2021** (2021).
46. J. Tariboon, S. K. Ntouyas, Quantum calculus on finite intervals and applications to impulsive difference equations, *Adv. Differ. Equ.*, **2013** (2013), 282.
47. M. Vivas-Cortez, M. A. Ali, A. Kashuri, I. B. Sial, Z. Zhang, Some new Newton's type integral inequalities for co-ordinated convex functions in quantum calculus, *Symmetry*, **12** (2020), 1476.
48. S. Zhao, S. I. Butt, W. Nazir, J. Nasir, M. Umar, Y. Liu, Some Hermite-Jensen-Mercer type inequalities for  $k$ -Caputo-fractional derivatives and related results, *Adv. Differ. Equ.*, **2020** (2020), 1–17.
49. J. Zhao, S. I. Butt, J. Nasir, Z. Wang, I. Tilli, Hermite-Jensen-Mercer type inequalities for Caputo fractional derivatives, *J. Funct. Spaces*, **2020** (2020), 7061549.



AIMS Press

©2021 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>)