

AIMS Mathematics, 9(1): 289–301. DOI: 10.3934/math.2024017 Received: 15 September 2023 Revised: 08 November 2023 Accepted: 14 November 2023 Published: 27 November 2023

http://www.aimspress.com/journal/Math

# Research article

# Extremal values of the first reformulated Zagreb index for molecular trees with application to octane isomers

Shabana Anwar<sup>1</sup>, Muhammad Kamran Jamil<sup>1</sup>, Amal S. Alali<sup>2,\*</sup>, Mehwish Zegham<sup>1</sup> and Aisha Javed<sup>3</sup>

- <sup>1</sup> Department of Mathematics, Riphah International University, Lahore, Pakistan
- <sup>2</sup> Department of Mathematical Sciences, College of Science, Princess Nourah bint Abdulrahman University, P.O. Box 84428, Riyadh 11671, Saudi Arabia
- <sup>3</sup> Abdus Salam School of Mathematical Sciences, Government College University, Lahore, Pakistan
- \* Correspondence: Email: asalali@pnu.edu.sa; ORCID: 0000-0001-7856-2861.

**Abstract:** A connected acyclic graph in which the degree of every vertex is at most four is called a molecular tree. A number associated with a molecular tree that can help to approximate the physical or chemical properties of the corresponding molecule is called a topological index. It is of great importance to investigate the relation between the structure and the thermodynamic properties of those molecules. In this paper, we investigated the extreme value of the first reformulated Zagreb index with a given order and degree of a graph. Further, we investigated the molecular trees that attain the maximum and minimum values. As an application, we presented the regression models to predict the acentric factor and entropy of octane isomers. Our extremal graphs give the minimum and the maximum acentric factor and entropy, which satisfied the experimental values.

**Keywords:** molecular tree; first reformulated Zagreb index; extremal trees; entropy; isomers **Mathematics Subject Classification:** 05C92, 05C90

# 1. Introduction

A graph is defined as  $\mathcal{J} = (\mathcal{V}_{\alpha}, \mathcal{E}_{\alpha})$ , where  $\mathcal{V}_{\alpha}$  is the graph's vertex set and  $\mathcal{E}_{\alpha}$  is its edge set, respectively. Structures of chemical compounds are represented by chemical bonds. One type of these bonds is a covalent bond in which the sharing of electrons takes place, and this sharing is indicated by lines called edges in chemical graphs and atoms as vertices. An edge between two vertices  $u_{\alpha}$  and  $v_{\alpha}$  is denoted by  $u_{\alpha}v_{\alpha}$ . The vertex degree denoted by  $\rho_{\mathfrak{J}}(u_{\alpha})$  of a vertex  $u_{\alpha}$  is measured by the total count of links connected with  $u_{\alpha}$ . The maximum degree in a graph is denoted as  $\Delta(\mathcal{J})$ , or simply  $\Delta$ . A single-component graph having no cycle with  $\triangle = 4$  is known as a molecular tree. The structures of acyclic molecules can be modeled by using molecular trees.

For a molecular graph, a topological descriptor is a numeral linked with this graph. This value can help to analyze some physical or chemical properties of the corresponding molecular structure. The family of Zagreb indices is an important family of topological indices, which contains various versions of the Zagreb indices. In 1972, the first and the second Zagreb indices were introduced [1,2].

The above two indices for  $\mathcal J$  are defined as:

$$M_{1}(\mathcal{J}) = \sum_{u_{\alpha} \in \mathcal{V}(\mathcal{J})} \rho(u_{\alpha})^{2},$$
$$M_{2}(\mathcal{J}) = \sum_{u_{\alpha}v_{\alpha} \in \mathcal{E}(\mathcal{G})} \rho(u_{\alpha})\rho(v_{\alpha}).$$

Miličević et al. [3] formulated the first and second Zagreb indices based on the edge degree. For the edge  $e = u_{\alpha}v_{\alpha}$ , the degree of e is  $\rho(e) = \rho(u_{\alpha}) + \rho(v_{\alpha}) - 2$  and the reformulated Zagreb descriptor is formulated by:

$$\begin{split} EM_1(\mathcal{J}) &= \sum_{e \in \mathcal{E}(\mathcal{J})} \rho(e)^2 = \sum_{u_\alpha v_\alpha \in \mathcal{E}(\mathcal{J})} \left( \rho(u_\alpha) + \rho(v_\alpha) - 2 \right)^2, \\ EM_2(\mathcal{J}) &= \sum_{e_1 \sim e_2} \rho(e_1) \rho(e_2). \end{split}$$

Extreme values of first and lower bounds for second reformulated Zagreb indices are calculated in [4]. A relation between a graph and its line graph is used to determine a relationship among original and reformulated Zagreb indices in [5]. In article [6], the authors discussed the maximum and minimum values of trees for  $EM_1(\mathcal{J})$  and  $EM_2(\mathcal{J})$ . In [7, 8], authors investigated the first, second, and third maximum Randić indices of molecular trees.

Ji et al. [9] investigated the external trees, unicyclic and bicyclic graphs with given order for the first reformulated Zagreb indices. Some important results for dendrimers, related to the above two indices are presented in [10]. In [11], authors found the extremal tricyclic graphs for the index. In [12, 13], the extremal values of molecular trees are calculated for Sombor indices. Bounds on  $EM_1(\mathcal{J})$  were observed by Milovanović et al. in [14], in terms of some graph parameters. The dependency of energy on cube of degree of vertices is shown in [15]. Neighborhood of Zagreb index on product of graphs is discussed in [16]. Su et al. [17] studied with given connectivity and determined the extremal graphs for the same index. The exact expressions for the same index as the first reformulated Zagreb index are presented in [18] for some familiar graph operations. For the general Randić index, the extremal molecular trees with the given number of pendant vertices are discussed in [19–21]. Liu and Li explored further properties of the molecular trees for the harmonic index in [22], and they also gave the extremal values of the harmonic descriptor of trees with  $\Delta = 4$ . Husin et al. [23] extended the results and determined the fourth maximum Randić index for the same class. Hundreds of topological indices have been introduced up to the date for history and latest results on this topic, we suggest [24–29].

In this paper we extended the work on the molecular trees for the first reformulated Zagreb index and determined the molecular trees, which give the extremal values of the index.

**AIMS Mathematics** 

2. Main results

In this section, we presented some auxiliary lemmas and our main results. In the following for  $1 \le i, j \le 4$ , the count of atoms having degree *i* is represented by  $n_i$ , while the count of bonds with one end atom having degree *i* and the other end atom containing degree *j* is represented by  $\zeta_{ij}$ .

For a molecular tree T we have the following information:

$$n_1 + n_2 + n_3 + n_4 = n, (2.1)$$

$$\zeta_{12} + \zeta_{13} + \zeta_{14} = n_1, \tag{2.2}$$

$$\zeta_{21} + 2\zeta_{22} + \zeta_{23} + \zeta_{24} = 2n_2, \tag{2.3}$$

$$\zeta_{31} + \zeta_{32} + 2\zeta_{33} + \zeta_{34} = 3n_3, \tag{2.4}$$

$$\zeta_{41} + \zeta_{42} + \zeta_{43} + 2\zeta_{44} = 4n_4, \tag{2.5}$$

$$n_1 + 2n_2 + 3n_3 + 4n_4 = 2(n-1).$$
(2.6)

Applying the definition of  $EM_1(T)$ ,

$$EM_{1}(T) = \sum_{u_{\alpha}v_{\alpha}\in\mathcal{E}(T)} (\rho(u_{\alpha}) + \rho(v_{\alpha}) - 2)^{2}$$
  
= $\zeta_{12} + 4\zeta_{13} + 9\zeta_{14} + 4\zeta_{22} + 9\zeta_{23} + 16\zeta_{24} + 16\zeta_{33} + 25\zeta_{34} + 36\zeta_{44}.$  (2.7)

From Eqs (2.1) and (2.6), we have

$$3n_1 + 2n_2 + n_3 = 2(1+n). (2.8)$$

Using Eqs (2.2)–(2.4) in (2.8), we get

$$\zeta_{14} = \frac{2n+2}{3} - \frac{4}{3}\zeta_{12} - \frac{10}{9}\zeta_{13} - \frac{2}{3}\zeta_{22} - \frac{4}{9}\zeta_{23} - \frac{1}{3}\zeta_{24} - \frac{2}{9}\zeta_{33} - \frac{1}{9}\zeta_{34}.$$

Solving Eqs (2.1) and (2.6), we have

$$n_2 + 2n_3 + 3n_4 = n - 2. \tag{2.9}$$

Using Eqs (2.3)–(2.5) in (2.9), we get

$$\frac{1}{2}(\zeta_{21}+2\zeta_{22}+\zeta_{23}+\zeta_{24}) + \frac{2}{3}(\zeta_{31}+\zeta_{32}+2\zeta_{33}+\zeta_{34}) + \frac{3}{4}(\zeta_{41}+\zeta_{42}+\zeta_{43}+2\zeta_{44}) = (n-2)$$
  
$$\zeta_{44} = \frac{n-5}{3} + \frac{1}{3}\zeta_{21} - \frac{1}{3}\zeta_{22} - \frac{5}{9}\zeta_{23} - \frac{2}{3}\zeta_{24} + \frac{1}{9}\zeta_{13} - \frac{7}{9}\zeta_{33} - \frac{8}{9}\zeta_{34}.$$

Substituting the values of  $\zeta_{14}$  and  $\zeta_{44}$  in Eq (2.7), we have

$$EM_1(T) = \zeta_{12} - 2\zeta_{13} - 14\zeta_{22} - 15\zeta_{23} - 11\zeta_{24} - 14\zeta_{33} - 8\zeta_{34} + 18n - 54.$$
(2.10)

The following result provides information about the maximum value of  $EM_1(T)$  for the molecular trees and the proof is given after some auxiliary lemmas.

AIMS Mathematics

Volume 9, Issue 1, 289-301.

**Theorem 2.1.** Let T be a molecular tree of order  $n \ge 5$ , then the maximum value of the first reformulated Zagreb index is

$$EM_1(T) \leq \begin{cases} 18n - 64, n \equiv 0 \pmod{3}, \\ 18n - 72, n \equiv 1 \pmod{3}, \\ 18n - 54, n \equiv 2 \pmod{3}, \end{cases}$$

and the maximum first reformulated Zagreb index is achieved if, and only if, one of the following conditions is satisfied:

- *T* has exactly one edge of degree one and one edge of degree sixteen,
- two edges are of degree four and degree of one edge is twenty five,
- degree of all the edges are either nine or thirty six.

Now we present some lemmas.

**Lemma 2.2.** Let a molecular tree T has at least two vertices of degree three, then T cannot attain the maximal first reformulated Zagreb Index.

*Proof.* Suppose we have a molecular tree *T* with at least two vertices of degree three, and  $u_{\alpha}$  and  $v_{\alpha}$  are three degree vertices, i.e.,  $\rho(u_{\alpha}) = \rho(v_{\alpha}) = 3$ . Based on adjacency of  $u_{\alpha}$  and  $v_{\alpha}$ , we have the following two cases.

**Case I.** When  $u_{\alpha}v_{\alpha} \notin \mathcal{E}(T)$ .

Suppose  $u_{\alpha_1}, u_{\alpha_2}, u_{\alpha_3}$  are the neighbors of  $u_{\alpha}$  in T and  $e_i = u_{\alpha}u_{\alpha_i}$  for  $1 \le i \le 3$ , and  $v_{\alpha_1}, v_{\alpha_2}, v_{\alpha_3}$  are the neighbors of  $v_{\alpha}$  in T and  $f_i = v_{\alpha}v_{\alpha_i}$  for  $1 \le i \le 3$ .

Further, suppose that  $\rho(u_{\alpha_1}) + \rho(u_{\alpha_2}) + \rho(u_{\alpha_3}) \le \rho(v_{\alpha_1}) + \rho(v_{\alpha_2}) + \rho(v_{\alpha_3})$ .

Now, we construct a new graph  $T' = T - u_{\alpha}u_{\alpha_2} + v_{\alpha}u_{\alpha_2}$ , and in T' we have  $d_{T'}(u_{\alpha}) = 2$  and  $d_{T'}(v_{\alpha}) = 4$ . Suppose  $A = \{e_1, e_2, e_3, f_1, f_2, f_3\}$ . Now, from the definition of T we have:

$$EM_{1}(T) - EM_{1}(T') = \{1 + \rho(u_{\alpha_{1}})\}^{2} + \{1 + \rho(u_{\alpha_{2}})\}^{2} + \{1 + \rho(u_{\alpha_{3}})\}^{2} + \{1 + \rho(v_{\alpha_{1}})\}^{2} + \{1 + \rho(v_{\alpha_{1}})\}^{2} - \{\rho(u_{\alpha_{1}})\}^{2} - \{\rho(u_{\alpha_{3}})\}^{2} - \{2 + \rho(u_{\alpha_{2}})\}^{2} - \{2 + \rho(v_{\alpha_{1}})\}^{2} - \{\rho(v_{\alpha_{2}}) + 2\}^{2} - \{\rho(v_{\alpha_{3}}) + 2\} < 0.$$

**Case II.** When  $u_{\alpha}v_{\alpha} \in \mathcal{E}(T)$ .

Here, we suppose that  $u_{\alpha_1} = u_{\alpha}$ ,  $v_{\alpha_1} = v_{\alpha}$  and  $e_1 = u_{\alpha}v_{\alpha} = f_1$ . Without loss of generality, take  $\rho(u_{\alpha_2}) + \rho(u_{\alpha_3}) \le \rho(v_{\alpha_2}) + \rho(v_{\alpha_3})$ . We construct a new tree  $T' = T - u_{\alpha}u_{\alpha_3} + v_{\alpha}u_{\alpha_3}$ . Let  $A = \{e_2, e_3, f_2, f_3\}$ , then

$$EM_{1}(T) - EM_{1}(T') = \{1 + \rho(u_{\alpha_{2}})\}^{2} + \{1 + \rho(u_{\alpha_{3}})\}^{2} + \{1 + \rho(v_{\alpha_{2}})\}^{2} + \{1 + \rho(v_{\alpha_{3}})\}^{2} - \{\rho(u_{\alpha_{2}})\}^{2} - \{1 + \rho(v_{\alpha_{2}})\}^{2} - \{2 + \rho(v_{\alpha_{3}})\}^{2} - \{2 + \rho(u_{\alpha_{3}})\}^{2} < 0,$$

which is a contradiction.

**Lemma 2.3.** If a molecular tree has at least two vertices of degree two, then it cannot attain the maximum reformulated Zagreb index.

**AIMS Mathematics** 

Volume 9, Issue 1, 289-301.

**Case I.** When  $u_{\alpha}v_{\alpha} \notin \mathcal{E}(T)$ .

Since  $\rho(u_{\alpha}) = 2 = \rho(v_{\alpha})$ , let  $e_1 = u_{\alpha}u_{\alpha_1}$ ,  $e_2 = u_{\alpha}u_{\alpha_2}$ ,  $f_1 = v_{\alpha}v_{\alpha_1}$  and  $f_2 = v_{\alpha}v_{\alpha_2}$ . Without loss of generality, suppose that  $\rho(u_{\alpha_1}) + \rho(u_{\alpha_2}) \le \rho(v_{\alpha_1}) + \rho(v_{\alpha_2})$ .

We construct a new molecular tree from T as  $T' = T - u_{\alpha}u_{\alpha_2} + v_{\alpha}u_{\alpha_2}$ , then

$$EM_{1}(T) - EM_{1}(T') = \rho(u_{\alpha_{1}})^{2} + \rho(u_{\alpha_{2}})^{2} + d(v_{\alpha_{1}})^{2} + \rho(v_{\alpha_{2}})^{2} - \{\rho(u_{\alpha_{1}}) - 1\}^{2} - \{1 + \rho(u_{\alpha_{2}})\}^{2} - \{1 + \rho(v_{\alpha_{1}})\}^{2} - \{1 + \rho(v_{\alpha_{2}})\}^{2} < 0.$$

**Case II.** When  $u_{\alpha}v_{\alpha} \in \mathcal{E}(T)$ .

Since the degree of  $u_{\alpha}$  and  $v_{\alpha}$  is two, suppose  $u_{\alpha_2}$  and  $v_{\alpha_2}$  are their other neighbors. The new molecular graph is constructed from *T* as  $T' = T - u_{\alpha}u_{\alpha_2} + v_{\alpha}u_{\alpha_2}$ . In this new graph, the degree of  $u_{\alpha}$  is one and the degree of  $v_{\alpha}$  is three.

$$EM_{1}(T) - EM_{1}(T') = \rho(u_{\alpha_{2}})^{2} + \rho(v_{\alpha_{2}})^{2} - \{1 + \rho(u_{\alpha_{2}})\}^{2} - \{1 + \rho(v_{\alpha_{2}})\}^{2} < 0.$$
  
$$EM_{1}(T) < EM_{1}(T').$$

In both cases we get a contradiction as we supposed T gives the maximum first reformulated Zagreb index.  $\Box$ 

**Lemma 2.4.** Let T be a molecular tree with at least one vertex of degree two, and at least one vertex of degree three, then T cannot attain the maximum first reformulated Zagreb index.

*Proof.* Let *T* be a molecular tree with a vertex  $u_{\alpha}$  of degree two and a vertex  $v_{\alpha}$  of degree three. **Case I.** Vertices  $u_{\alpha}$  and  $v_{\alpha}$  are not adjacent.

Since the degrees of  $u_{\alpha}$  and  $v_{\alpha}$  are two and three, respectively, let  $e_1 = u_{\alpha}u_{\alpha_1}, e_2 = u_{\alpha}u_{\alpha_2}, f_1 = v_{\alpha}v_{\alpha_1}, f_2 = v_{\alpha}v_{\alpha_2}$  and  $f_3 = v_{\alpha}v_{\alpha_3}$ . By deleting edge  $u_{\alpha}u_{\alpha_2}$  and adding an edge  $v_{\alpha}u_{\alpha_2}$  from *T*, we get another molecular graph *T'*.

$$EM_{1}(T) - EM_{1}(T') = \rho(u_{\alpha_{1}})^{2} + \rho(u_{\alpha_{2}})^{2} + \{1 + \rho(v_{\alpha_{1}})\}^{2} + \{1 + \rho(v_{\alpha_{2}})\}^{2} + \{1 + \rho(v_{\alpha_{3}})\}^{2} - \{\rho(u_{\alpha_{1}}) - 1\}^{2} - \{2 + \rho(v_{\alpha_{1}})\}^{2} - \{2 + \rho(v_{\alpha_{2}})\}^{2} - \{2 + \rho(v_{\alpha_{3}})\}^{2} \{2 + \rho(u_{\alpha_{2}})\}^{2} < 0.$$

**Case II.** Vertices  $u_{\alpha}$  and  $v_{\alpha}$  are adjacent.

Let the vertex  $u_{\alpha_2}$  be different from  $v_{\alpha}$  as adjacent to  $u_{\alpha}$ , and vertices  $v_{\alpha_1}$  and  $v_{\alpha_2}$  different from  $u_{\alpha}$  are adjacent the vertex  $v_{\alpha}$ . The new molecular graph T' is constructed from T by deleting the edge  $u_{\alpha}u_{\alpha_2}$ and adding an edge  $v_{\alpha}u_{\alpha_2}$ . When  $u_{\alpha}$  and  $v_{\alpha}$  are neighbors of each other, then  $u_{\alpha_1}$  and  $v_{\alpha_1}$  are actually vertices  $u_{\alpha}$  and  $v_{\alpha}$ . Graph T' is obtained by deleting edge  $u_{\alpha}u_{\alpha_2}$  and adding edge  $v_{\alpha}u_{\alpha_2}$ .

$$EM_{1}(T) - EM_{1}(T') = \{\rho(u_{\alpha_{2}})\}^{2} + \{1 + \rho(v_{\alpha_{2}})\}^{2} + \{\rho(v_{\alpha_{3}}) + 1\}^{2} - \{\rho(v_{\alpha_{2}}) + 2\}^{2} - \{\rho(v_{\alpha_{3}}) + 2\}^{2} - \{\rho(u_{\alpha_{2}}) + 2\}^{2} < 0.$$

From the above three Lemmas, we draw following conclusion.

**AIMS Mathematics** 

Volume 9, Issue 1, 289–301.

#### 2.1. Conclusion

If T is a molecular tree with the maximum first reformulated Zagreb index, then T must satisfy one of the following three conditions.

- all vertices of *T* have a degree of either one or four,
- exactly one vertex of the graph is of degree two and remaining vertices are of degree one or four,
- exactly one vertex of the graph is of degree three and remaining vertices are of degree one or four.

#### 2.2. Proof of Theorem 2.1

*Proof.* For any molecular tree T, we have the following formula for the first reformulated Zagreb index derived in Eq (2.10)

$$EM_1(T) = 18n - 54 + \zeta_{12} - 2\zeta_{13} - 14\zeta_{22} - 15\zeta_{23} - 11\zeta_{24} - 14\zeta_{33} - 8\zeta_{34}.$$

Let T be a molecular tree with the maximum value of the first reformulated Zagreb index. Based on the order of T, we have the three cases given as:

**Case I.** When  $n \equiv 0 \mod (3)$ .

Let  $n = 3\hbar$  and  $\hbar \ge 2$ . From Eq (2.9), we get

$$n_2 + 2n_3 + 3n_4 = 3\hbar - 2. \tag{2.11}$$

Since *T* has the maximal first reformulated Zagreb index, then by Lemmas 2.2–2.4, the solution of the Eq (2.11) exists if, and only if,  $n_4 = \hbar - 1$ ,  $n_3 = 0$  and  $n_2 = 1$ . Putting these values in Eq (2.10), we have,

$$EM_1(T) = 18n - 54 + \zeta_{12} - 11\zeta_{24}.$$

Moreover,

$$EM_1(T) = \begin{cases} 18n - 76, & if \zeta_{12} = 0, \zeta_{24} = 2, \\ 18n - 64, & if \zeta_{12} = 1 = \zeta_{24}. \end{cases}$$

However,

$$EM_1(T) \le 18n - 64,$$

gives the maximum value and the equality holds if, and only if, T has exactly one edge of degree one and one vertex edge of degree sixteen.

**Case II.** When  $n \equiv 1 \mod (3)$ . Let  $n = 3\hbar + 1$ , and  $\hbar \ge 2$ , then Eq (2.9) becomes

$$n_2 + 2n_3 + 3n_4 = 3\hbar - 1. \tag{2.12}$$

From Lemmas 2.2–2.4, the solution of Eq (2.12) exists if, and only if,  $n_4 = \hbar - 1$ ,  $n_2 = 0$  and  $n_3 = 1$ . In the case of all  $\zeta_{ij} = 0$ , if any of i = 2 or j = 2, then  $\zeta_{33}=0$ . Thus Eq (2.10) reduces to,

$$EM_1(T) = 18n - 54 - 2\zeta_{13} - 8\zeta_{34}.$$

AIMS Mathematics

Volume 9, Issue 1, 289-301.

Moreover,

$$EM_{1}(T) = \begin{cases} 18n - 78, & if\zeta_{34} = 3, \zeta_{13} = 0, \\ 18n - 66, & if\zeta_{13} = 2, \zeta_{34} = 1. \end{cases}$$

Hence,

# $EM_1(T) \le 18n - 66,$

and the equality holds if, and only if, the degree of two edges of the molecular tree are four and the degree of one edge is twenty five.

#### **Case III.** When $n \equiv 2 \mod (3)$ .

Let  $n = 3\hbar + 2$  for  $\hbar \ge 2$ , and from Eq (2.9) we have  $n_2 + 2n_3 + 3n_4 = 3\hbar$ . As in previous cases, the solution of the equation exists if, and only if,  $n_4 = \hbar$ ,  $n_3 = 0$  and  $n_2 = 0$ , then  $\zeta_{ij} = 0$ , for i = 2, 3 or j = 2, 3. Hence, the first reformulated Zagreb index is

$$EM_1(T) = 18n - 54$$

and the above is attained if, and only if, the degree of all the edges are either nine or thirty six, which completes the proof.  $\Box$ 

#### 3. Minimum value of the reformulated Zagreb index

In the following, we will work on the minimal first reformulated Zagreb index of molecular trees for  $2 \le \Delta \le 4$ .

**Lemma 3.1.** Let T be a molecular tree with  $\triangle = 3$  and with two vertices of degree three, then T cannot attain the minimal first reformulated Zagreb index.

*Proof.* Let *T* be a molecular tree on *n* vertices and  $\triangle = 3$  with two vertices of degree three, i.e.,  $\rho(u_{\alpha}) = \rho(v_{\alpha}) = 3$  and with the minimal first reformulated Zagreb index. Suppose that there is a vertex  $w_{\alpha}$  such that  $\rho(w_{\alpha}) = 1$ . Let  $u_{\alpha 1}, u_{\alpha 2}, u_{\alpha 3}$  be neighbors of  $u_{\alpha}$  and  $w_{\alpha 1}$  is the neighbor of  $w_{\alpha}$  in *T*. Let's construct a new graph *T'* as,  $T' = T - u_{\alpha}u_{\alpha 1} + w_{\alpha}u_{\alpha 1}$ . Further we have,

$$EM_1(T) - EM_1(T') = 2(\rho(u_{\alpha_1}) + \rho(u_{\alpha_2}) + \rho(u_{\alpha_3})) - 2\rho(w_{\alpha_1}) + 4 > 0.$$

This result shows that the molecular tree is not minimal for two vertices of degree three.

Following lemmas can be proved similarly.

**Lemma 3.2.** Let T be a molecular tree with maximum degree four and two vertices of degree four, then the first reformulated Zagreb index of T is not minimum.

**Lemma 3.3.** Let *T* be a molecular tree of order *n* having a maximum degree four with one vertex of degree four and one vertex of degree three, then the first reformulated Zagreb descriptor cannot be minimum for *T*.

*Proof.* Let *T* be a tree with  $\triangle = 3$  and  $\rho(u_{\alpha}) = 3$  and  $\rho(w_{\alpha}) = 4$ . Suppose that  $u_{\alpha}$  has neighbors  $u_{\alpha_1}, u_{\alpha_2}$  and  $u_{\alpha_3}$  and  $w_{\alpha}$  has neighbor  $w_{\alpha_1}$ . Now, construct *T'* such that  $T' = T - u_{\alpha}u_{\alpha_1} + w_{\alpha}u_{\alpha_1}$ , then  $EM_1(T) - EM_1(T') > 0$ .

AIMS Mathematics

Volume 9, Issue 1, 289–301.

**Theorem 3.4.** Let a molecular tree with n number of vertices such that  $n \ge 5$  and  $2 \le \Delta \le 4$ , then the minimal value of the first reformulated Zagreb index is given as

$$EM_{1}(T) \geq \begin{cases} n-10 \text{ for } \triangle = 2, \text{ equality holds when } T \text{ is a path,} \\ 4n-2 \text{ for } \triangle = 3, \text{ equality holds when } T \text{ contains two edges of degree four,} \\ 4n+20 \text{ for } \triangle = 4, \text{ equality holds when } T \text{ contains three edges of degree nine} \end{cases}$$

*Proof.* • If  $\Delta = 2$ , then clearly tree is a path graph and we have  $EM_1(T) = 4n - 10$ .

• For  $\Delta = 3$ . By Lemma 3.1, we have only two molecular trees:

(1) one edge of degree four, denotes it  $T_1$ ,

(2) two edges of degree four, denotes it by  $T_2$ .

Further,  $EM_1(T_1) = 4n$  and  $EM_1(T_2) = 4n - 2$ , clearly  $EM_1(T_1) > EM_1(T_2)$ .

• For  $\Delta = 4$ . By Lemmas 3.2 and 3.3, we have the following molecular trees:

(1) two edges of degree nine, denotes it  $T_3$ ,

(2) three edges of degree nine, denotes it by  $T_4$ .

Further,  $EM_1(T_3) = 4n + 24$  and  $EM_1(T_4) = 4n + 20$ , and clearly  $EM_1(T_3) > EM_1(T_4)$ , which completes the proof.

## 4. Application to octane isomers

In this section, we will present a possible application of our work. The acentric factor is a measure of the non-centricity of molecules. As the acentric factor increases, the vapor pressure goes down, resulting in higher boiling points [30, 31]. Entropy is a measure of the unavailability of a system's energy to do work. It is a measure of disorder. The greater the disorder in a molecule, the greater the entropy [32, 33]. Here, 18 octane isomers [34, 35] with their acentric factors and entropy measure are considered, as well as the first reformulated Zagreb indices of these octane isomers. This information is provided in Table 1. Several papers have been written on the correlation between properties and topological indices of molecular graphs, and we refer [12, 15] and references therein. A linear regression through Microsoft Excel is performed on the data in Table 1. We obtain the following linear equations that can predict the acentric factor and entropy of octane isomers. Further, from Table 1 we can notice that the 2,2,3,3,-tetramethyl butane (18th isomer) has the maximum reformulated Zagreb index with the minimum acentric factor and entropy. In Theorem 2.1, we can notice that the same molecular tree attained the maximum value. This implies that the chemical trees with the maximum reformulated Zagreb index have the minimum acentric factor and entropy measure.

Acentric factor =  $-0.0021 \times RM_1 + 0.4351$ , Entropy =  $-0.258 \times RM_1 + 117.77$ ,

with the correlation coefficients (R) 0.9841 and 0.9608, respectively. These results are comparable with the models already discussed in the literature. Figures 1 and 2 show the close relation between the acentric factor and entropy and the reformulated Zagreb index. These results imply that the reformulated Zagreb index can be a good predictor of the acentric factor and entropy for octane isomers.



**Figure 1.** The relationship between acentric factor and the reformulated Zagreb index for 18 isomers.



Figure 2. The relationship between entropy and the reformulated Zagreb index for 18 octane isomers.

Volume 9, Issue 1, 289–301.

No.	Isomer	Acent. Fac.	Entropy	$EM_1$
1	• • • • • • • •	0.397898	111.67	22
2		0.377916	109.84	30
3		0.371002	111.26	32
4		0.371504	109.32	32
5		0.362472	109.43	34
6		0.339426	103.42	52
7		0.348247	108.02	42
8		0.344223	106.98	40
9		0.356830	105.72	38
10		0.322596	104.74	56
11		0.340345	106.59	44
12	- <u>I</u>	0.332433	106.06	44
13		0.306899	101.48	60
14		0.300816	101.31	66
15		0.305370	104.09	60
16		0.293177	102.06	68
17		0.317422	102.39	52
18		0.255294	93.06	90

**Table 1.** Acentric factor, entropy and the reformulated Zagreb index values of 18 octane isomers.

## 5. Conclusions

In the field of chemical graph theory, topological indices play an important role. Topological indices are numerical values that are used to correlate the chemical structure of molecules by a graph with

chemical properties. Many distance-based and degree-based topological indices have been introduced. Among all existing topological indices, degree-based indices are very helpful in hydrocarbons, in the field of pharmacy and the preparation of drugs used against diseases like cancer. In this article, we discussed the behavior of the first reformulated Zagreb index for molecular trees, which is the generalized form of of the first Zagreb index. Here, we calculated the first reformulated Zagreb index for molecular trees of order n and then calculated the results for extremal (maximal, minimal) values. The results obtained can be useful in extracting data for molecular trees and in discussing many properties of these molecular tree graphs, like melting point, boiling point, latent heat of fusion, entropy, enthalpy etc. In the end, we presented an application of the first reformulated Zagreb index, acentric factor and entropy graphically.

# Use of AI tools declaration

The authors declare they have not used Artificial Intelligence (AI) tools in the creation of this article.

## Acknowledgments

The authors extend their appreciation to Princess Nourah bint Abdulrahman University for funding this research under Researchers Supporting Project number (PNURSP2023R231), Princess Nourah bint Abdulrahman University, Riyadh, Saudi Arabia.

## **Conflict of interest**

Authors declare there is no conflict of interest in publishing the article.

## References

- I. Gutman, T. Nenad, Graph theory and molecular orbitals. Total φ-electron energy of alternant hydrocarbons, *Chem. Phys. Lett.*, **17** (1972), 535–538. https://doi.org/10.1016/0009-2614(72)85099-1
- 2. S. Nikolić, G. Kovačević, A. Miličević, N. Trinajstić, The Zagreb indices 30 years after, *Croat. Chem. Acta*, **76** (2003), 113–124.
- 3. A. Miličević, S. Nikolić, N. Trinajstić, On reformulated Zagreb indices, *Mol. Divers.*, **8** (2004), 393–399. https://doi.org/10.1023/B:MODI.0000047504.14261.2a
- 4. A. Ilić, B. Zhou, On reformulated Zagreb indices, *Discrete Appl. Math.*, **160** (2012), 204–209. https://doi.org/10.1016/j.dam.2011.09.021
- 5. A. Miličević, S. Nikolić, N. Trinajstić, On reformulated Zagreb indices, *Mol. Divers.*, **8** (2004), 393–399. https://doi.org/10.1023/B:MODI.0000047504.14261.2a
- 6. A. Ghalavand, A. R. Ashrafi, Extremal trees with respect to the first and second reformulated Zagreb index, *Malaya J. Mate.*, **5** (2017), 524–530. https://doi.org/10.26637/mjm503/006

- I. Gutman, O. Miljković, G. Caporossi, P. Hansen, Alkanes with small and large Randić connectivity indices, *Chemi. Phys. Lett.*, **306** (1999), 366–372. https://doi.org/10.1016/S0009-2614(99)00472-8
- 8. I. Gutman, O. Miljković, Molecules with smallest connecivity tivity indices, *Match-Commun. Math. Co.*, **41** (2000), 57–70.
- 9. S. Ji, X. Li, B. Huo, On reformulated Zagreb indices with respect to acyclic, unicyclic and bicyclic graphs, *Match-Commun. Math. Co.*, **72** (2014), 723–732.
- 10. N. De, Reformulated Zagreb indices of dendrimers, Math. Aeterna, 3 (2013), 133–138.
- 11. S. Ji, Y. Qu, X. Li, The reformulated Zagreb indcies of tricyclic graphs, *Appl. Math. Comput.*, **268** (2015), 590–595. https://doi.org/10.1016/j.amc.2015.06.058
- 12. H. Deng, Z. Tang, R. Wu, Molecular trees with extremal values of Sombor indices, *Int. J. Quantum. Chem.*, 2021. https://doi.org/10.1002/qua.26622
- M. Chemometrics, Molecular Descriptors: The free online resource, Milano Chemometrics and QSAR Research Group, 2017. Available from: http://www.moleculardescriptors.eu/ dataset/dataset.htm.
- E. I. Milovanović, I. Z. Milovanović, E. C. Dolićanin, E. Glogić, A note on the first reformulated Zagreb index, *Appl. Math. Comput.*, 273 (2016), 16–20.
- 15. B. Furtula, I. Gutman, A forgotten topological index, J. Math. Chem., 53 (2015), 1184–1190. https://doi.org/10.1007/s10910-015-0480-z
- 16. S. Mondal, N. De, A. Pal, On neighborhood Zagreb index of product graphs, *J. Mol. Struct.*, 2021. https://doi.org/10.1016/j.molstruc.2020.129210
- 17. G. Su, L. Xiong, L. Xu, B. Ma, On the maximum and minimum first reformulated Zagreb index of graphs with connectivity at most k, *Filomat*, 25 (2011), 75–83. https://doi.org/10.2298/FIL1104075S
- 18. M. K. Jamil, I. Tomescu, First reformulated Zagreb index and some graph operations, *Ars Combinatoria*, **138** (2018), 193–209.
- 19. X. Li, Y. Yang, Best lower and upper bounds for the Randic index  $R_{-1}$  of chemical trees, *Match-Commun. Math. Co.*, **52** (2004), 147–156.
- 20. X. Li, Y. Shi, L. Zhong, Minimum general Randić index on chemical trees with given order and number of pendent vertices, *Match-Commun. Math. Co.*, **60** (2008), 539–554.
- R. Xing, B. Zhou, N. Trinajstić, Sum-connectivity index of molecular trees, J. Math. Chem., 48 (2010), 583–591. https://doi.org/10.1007/s10910-010-9693-3
- 22. S. Liu, J. Li, Some properties on the harmonic index of molecular trees, *ISRN Appl. Math.*, **2014** (2014). http://dx.doi.org/10.1155/2014/781648
- 23. N. H. M. Husin, R. Hasni, Z. Du, A. Ali, More results on extremum Randić indices of (molecular) trees, *Filomat*, 2 (2018), 3581–3590. https://doi.org/10.2298/FIL1810581H
- M. Imran, R. Ismail, M. Azeem, M. K. Jamil, E. H. A. Al-Sabri, Sombor topological indices for different nanostructures, *Heliyon*, 2023. https://doi.org/10.1016/j.heliyon.2023.e20600

- M. Waheed, U. Saleem, A. Javed, M. K. Jamil, Computational aspects of entropy measures for metal organic frameworks, *Mol. Phys.*, 2023. https://doi.org/10.1080/00268976.2023.2254418
- 26. R. Luo, K. Dawood, M. K. Jamil, M. Azeem, Some new results on the face index of certain polycyclic chemical networks, *Math. Biosci. Eng.*, 20 (2023), 8031–8048. https://doi.org/10.3934/mbe.2023348
- 27. I. Gutman, I. Redžepović, B. Furtula, On the product of Sombor and modified Sombor indices, *Open J. Discrete Appl. Math.*, **6** (2023), 1–6. https://doi.org/10.30538/psrp-odam2023.0083
- 28. H. Liu, Extremal (n,m)-graphs with respect to VDB topological indices, *Appl. Math.*, **6** (2023), 16–20. https://doi.org/10.3934/mbe.2023240
- 29. Y. Yao, The smallest sum-connectivity index on trees with n vertices and k pendant vertices, *Open J. Discrete Appl. Math.*, **2** (2019), 23–30. https://doi.org/10.30538/psrp-odam2019.0013
- D. R. Schreiber, S. P. Kenneth, Equation of state in the acentric factor system, *Fluid Phase Equilibr.*, 46 (1989), 113–130. https://doi.org/10.1016/0378-3812(89)80032-9
- 31. G. M. Kontogeorgis, D. P. Tassios, Critical constants and acentric factors for long-chain alkanes suitable for corresponding states applications, *Chem. Eng. J.*, **66** (1997), 35–49. https://doi.org/10.1016/S1385-8947(96)03146-4
- 32. A. Rényi, *On measures of entropy and information*, Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability, Volume 1: Contributions to the Theory of Statistics, University of California Press, **4** (1961), 547–562.
- C. Ricotta, L. Szeidl, Diversity partitioning of Rao's quadratic entropy, *Theor. Popul. Biol.*, 76 (2009), 299–302. https://doi.org/10.1016/j.tpb.2009.10.001
- 34. C. Tamilarasi, The pivotal role of degree-based and neighborhood degree-sum-based topological indices in predicting the Physico-Chemical properties of N-octane isomers, *Ann. Rom. Soc. Cell Biol.*, 25 (2021), 4481–4489.
- 35. B. A. Ravindran, D. Togarichetu, V. M. Gangadharaiah, L. Veerabhadraiah, QSPR analysis of VL index with octane isomers, *AIP Conf. Proc.*, **2649** (2023). https://doi.org/10.1063/5.0117457



© 2024 the Author(s), licensee AIMS Press. This is an open access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0)